



MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Lumbar Fusion for Adults
Policy Number	7.01.90
Category	Technology Assessment
Original Effective Date	02/19/15
Committee Approval Date	01/21/16, 01/19/17, 01/18/18, 06/21/18, 12/20/18, 07/18/19, 01/16/20, 02/18/21, 02/17/22
Revised Effective Date	02/17/22
Archived Date	N/A
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit. • If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, lumbar spinal fusion with decompression in adult patients has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following criteria have been met:
 - A. The patient is a candidate for lumbar decompression (*refer to Corporate Medical Policy #7.01.97 Lumbar Decompression*).
 - B. The procedure is performed for actual or anticipated iatrogenic instability from decompression when **EITHER** of the following criteria is met:
 1. actual or anticipated instability is identified intra-operatively, created by disruption of the posterior elements due to facet joint excision that exceeds 50% bilaterally or 75% or more of a single facet during spinal decompression; or
 2. confirmatory imaging includes **ANY** of the following (not required when instability is created and/or identified intra-operatively):
 - a. post-operative instability created by the disruption of the posterior elements due to facet joint excision that exceeds 50% bilaterally or 75% or more of a single facet; or
 - b. removal of the pars interarticularis is performed that requires fusion to stabilize; or
 - c. pars fracture; or
 - d. previous spinal decompression that resulted in iatrogenic spondylolisthesis; or
 - e. Meyerding Grade I or higher spondylolisthesis with foraminal stenosis.
 - C. The patient has no untreated, underlying behavioral health conditions/issues (e.g., depression, chronic pain syndrome, secondary gain, opioid use or alcohol use disorder) as a contributor to chronic pain.
- II. Based upon our criteria and assessment of the peer-reviewed literature, lumbar fusion (arthrodesis) without decompression in adult patients has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:
 - A. The patient has a significant level of pain on a daily basis, defined as **EITHER** of the following:

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 2 of 11

1. Visual Analog Scale (VAS) /Numeric Rating Scale (NRS) greater than or equal to seven on a daily basis; or
2. severe, disabling, crippling, or incapacitating pain.
- B. The patient has clinically significant functional impairment (e.g., inability to perform household chores, prolonged standing or essential job functions).
- C. The patient has not experienced clinically meaningful improvement with **EITHER** of the following for at least three consecutive months, unless contraindicated, except for discogenic lower back/degenerative disc disease:
 1. prescription-strength analgesics, steroids, and/or NSAIDs;
 2. provider-directed exercise program prescribed by a physical therapist, chiropractic provider, or osteopathic or allopathic physician.
- D. The patient has no untreated, underlying behavioral health conditions/issues (e.g., depression, chronic pain syndrome, secondary gain, opioid use or alcohol use disorder) as a contributor to chronic pain.
- E. The procedure is performed for **ANY** of the following conditions:
 1. spondylolisthesis, with spondylolysis confirmed by imaging with **ANY** of the following:
 - a. multi-level spondylolysis on plain x-rays;
 - b. symptomatic Meyerding Grade 1 or 2 spondylolisthesis (anterolisthesis) with plain x-rays supporting progression of anterolisthesis;
 - c. symptomatic Meyerding Grade 3 or higher spondylolisthesis (anterolisthesis) demonstrated on plain x-rays with 50% or more anterior slippage **OR** plain x-rays supporting progression of anterolisthesis; or
 - d. progressive spinal pain with confirmatory imaging of progression of spondylolisthesis;
 2. degenerative spondylolisthesis without spondylolysis, when confirmatory imaging results show that **EITHER** the following is present:
 - a. dynamic segmental instability documented by flexion-extension plain x-rays **OR** comparison of a supine and upright image, with a difference in translational alignment between vertebrae greater than three mm between views; or
 - b. Meyerding grade II or higher spondylolisthesis (i.e., instability);
 3. discogenic lower back/degenerative disc disease, when **ALL** of the following are met:
 - a. presence of chronic, unremitting, discogenic axial lower back pain and associated disability secondary to single-level degenerative lumbar disc disease (DDD) for at least one year;
 - b. structured, physician-supervised, multi-modal, non-operative management of medical care with licensed healthcare professionals, which includes regularly scheduled appointments, follow-up evaluation, and less than clinically meaningful improvement with at least **TWO** of the following for at least 12 consecutive months, unless contraindicated:
 - i. prescription-strength analgesics, steroids, and/or NSAIDs;
 - ii. provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician;
 - iii. epidural steroid injection(s)/selective nerve root blocks; or
 - iv. facet joint injection(s)/medial branch block(s)/radiofrequency ablation(s); and
 - c. moderate-to-severe single-level disc degeneration has been confirmed on plain x-rays and advanced diagnostic imaging studies (i.e., CT, MRI);
 4. initial disc herniation, when **BOTH** of the following criteria are met:
 - a. Patient is a candidate for initial primary lumbar discectomy; and
 - b. Any of the following are present:
 - i. primary extraforaminal disc herniation at L5-S1, in which a far lateral approach is not feasible because of the presence of the iliac wings;
 - ii. primary foraminal disc herniation for which facet resection is necessary to retrieve the disc, which will result in iatrogenic instability; or
 - iii. primary disc herniation in the lumbar spine that is at the level of the spinal cord (i.e., low lying conus medullaris);
 5. recurrent disc herniation, when **ALL** of the following criteria are met:
 - a. The patient is a candidate for repeat lumbar discectomy (*refer to Corporate Medical Policy #7.01.98 Lumbar Microdiscectomy*); and

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 3 of 11

- b. Confirmatory plain x-rays demonstrate neural structure compression, and
 - c. Plain x-ray evidence of anterolisthesis is present, resulting in **EITHER** of the following:
 - i. segmental instability with a minimum of three mm or more of anterior translation displacement of the vertebra on the adjacent vertebra below; or
 - ii. Meyerding grade 2 or higher spondylolisthesis (i.e., instability);
 - 6. second or greater recurrent disc herniation when the patient is a candidate for repeat lumbar discectomy; and
 - 7. isthmic spondylolisthesis, when congenital or acquired pars defect is documented by imaging studies.
- III. Based upon our criteria and assessment of the peer-reviewed literature, repeat lumbar fusion (arthrodesis) at the same level in adult patients has been medically proven to be effective and, therefore, is considered **medically appropriate** when **EITHER** of the following criteria is met:
 - A. Post-operative plain x-rays show evidence of malposition or implant failure (e.g., migration, pedicle screw breakage, pedicle screw loosening, dislodged hooks, rod breakage, rod bending, rod loosening, loss of curve correction, decompensation, etc.).
 - B. **ALL** of the following criteria are met:
 - 1. The patient meets criteria for lumbar fusion (*see Policy Statement I or II*);
 - 2. More than six months have elapsed since the last fusion (arthrodesis) surgery; and
 - 3. Post-operative confirmatory imaging documenting pseudarthrosis includes ANY of the following:
 - a. magnetic resonance imaging (MRI);
 - b. computed tomography (CT); or
 - c. plain x-rays.
- IV. Based upon our criteria and assessment of the peer-reviewed literature, lumbar fusion (arthrodesis) for adjacent segment degeneration in adult patients has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following are met:
 - A. The patient meets criteria for lumbar fusion (*see Policy Statement I or II*).
 - B. The prior lumbar fusion (arthrodesis) procedure at an adjacent level was performed at least six months prior.
 - C. Evidence is present of anterolisthesis on plain x-rays resulting in BOTH of the following:
 - 1. segmental instability with three mm or more of anterior translation displacement of the vertebra on the adjacent vertebra below; and
 - 2. Meyerding grade II or higher spondylolisthesis (i.e., instability).
 - D. Neural structure compression is demonstrated by plain x-rays.
 - E. The patient experienced significant initial relief of symptoms following prior spinal fusion(s).
- V. Based upon our criteria and assessment of the peer-reviewed literature, lumbar discectomy and fusion following failed lumbar disc arthroplasty implant in adult patients has been medically proven to be effective and, therefore, is considered **medically appropriate** when **EITHER** of the following criteria is met:
 - A. Post-operative plain x-rays show evidence of implant malposition or implant failure (e.g., subsidence, loosening, infection, dislocation/subluxation, vertebral body fracture, dislodgement).
 - B. **ALL** of the following are met;
 - 1. Patient meets criteria for lumbar fusion (*see Policy Statement I or II*);
 - 2. Post-operative MRI /CT confirms evidence of neural structure compression (e.g., either retained disc material or a recurrent disc herniation); and
 - 3. More than six months have elapsed since disc arthroplasty surgery.
- VI. Based upon our criteria and assessment of the peer-reviewed literature, lumbar spinal fusion has not been medically proven to be effective and is, therefore, considered **not medically necessary** for **ANY** of the following sole indications:
 - A. disc herniation in the absence of **ANY** of the following:
 - 1. primary extraforaminal disc herniation at L5-S1, in which a far lateral approach is not feasible because of the presence of the iliac wings;
 - 2. primary foraminal disc herniation for which facet resection is necessary to retrieve the disc, which will result in iatrogenic instability; or

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 4 of 11

3. primary disc herniation in the lumbar spine that is at the level of the spinal cord (i.e., low-lying conus medullaris).
 - B. multi-level degenerative disc disease without instability;
 - C. neuro-compressive pathology;
 - D. facet joint disorders without instability;
 - E. initial discectomy/laminectomy without instability;
 - F. spondylolysis without spondylolisthesis; or
 - G. as an adjunct to primary decompression of central and/or lateral recess stenosis, in the absence of instability, spondylolisthesis, or an actual or anticipated bony resection that will result in iatrogenic instability.
- VII. Based upon our criteria and assessment of the peer-reviewed literature, the following devices/procedures have not been medically proven to be effective and, therefore, are considered **investigational** under circumstances that include, but are not limited to, the following:
- A. The procedure involves minimally invasive surgical approaches using only indirect visualization (e.g., endoscopic fusion, percutaneous fusion (video-imaging)).
 - B. The device/implant has not been approved by the U.S. Food and Drug Administration (FDA).
 - C. The procedure involves a dynamic (intervertebral) stabilization device (e.g., Dynesys, Stabilimax NZ).

Refer to Corporate Medical Policy #7.01.83 Minimally Invasive/Minimal Access Techniques for Lumbar Interbody Fusion.

Refer to Corporate Medical Policy #7.01.97 Lumbar Decompression.

Refer to Corporate Medical Policy #7.01.98 Lumbar Microdiscectomy.

Refer to Corporate Medical Policy #11.01.03 Experimental and Investigational Services.

POLICY GUIDELINES

- I. Documentation of nicotine-free status is required, established by **EITHER** of the following:
 - A. Patient is a never-smoker.
 - B. Patient has refrained from tobacco use and/or nicotine replacement therapy for at least six weeks prior to the planned spinal fusion surgery, as evidenced by lab results (cotinine level of less than or equal to 10ng/mL).
- II. The following are the minimum documentation requirements needed to complete a spinal surgery prior authorization request:
 - A. CPT codes, ICD-10 codes, and disc levels or motion segments involved for planned surgery must be provided.
 - B. Detailed documentation must also be provided of the type of provider-directed conservative treatment (e.g., interventional pain management procedures/injections, medication, physical therapy, chiropractic, or other provider-directed active exercise program) that includes response to conservative treatment, as well as the most recent imaging reports performed, read, and interpreted by an independent radiologist whose report shall supersede any discrepancies (when present) in interpretation.
 - C. The submission must include standing flexion-extension plain x-rays for spinal fusion surgery requests based upon indications of instability.
- III. **URGENT/EMERGENT CONDITIONS**: All patients being evaluated for spine surgery should be screened for indications of a medical condition that requires urgent/emergent treatment. The presence of such indications/condition warrants definitive surgical treatment. Provider-directed, non-surgical management, proof of smoking cessation, and absence of unmanaged significant behavioral health disorders are NOT required. Confirmatory advanced imaging, such as a CT scan or MRI, is required. Urgent/emergent conditions for thoracolumbar fusion (arthrodesis) include **ANY** of the following:
 - A. traumatic spinal fractures or dislocations, with or without neural compression, when instability is present or decompression of the spinal canal is anticipated to result in instability;
 - B. infection (e.g., discitis, epidural abscess, osteomyelitis) when instability is present or debridement and/or decompression is anticipated to result in instability;

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 5 of 11

- C. primary or metastatic neoplastic disease causing pathologic fracture, cord compression when instability is present or resection and/or decompression is anticipated to result in instability;
- D. congenital, neuromuscular, or infantile/juvenile/adolescent idiopathic scoliosis; or
- E. documentation of severe debilitating pain and/or dysfunction to the point of being incapacitated.

DESCRIPTION

Low-back pain affects approximately 90% of the U.S. population at some point in their lives and may be caused by a wide variety of conditions. Conservative management typically consists of rest, exercise, analgesics, local injections, lumbar bracing, physical therapy, and chiropractic care. Generally, conservative therapy is not recommended in the presence of progressive neurological deficits, when spinal fracture or dislocation is unstable or for progressive spinal deformity. When conservative management is attempted and fails, surgery may be required for conditions with underlying pathology as determined by radiological findings.

Spinal fusion/arthrodesis, also known as spondylodesis or spondylosynesis, is a well-established surgical technique for infectious conditions of the spine (e.g., spinal tuberculosis). It has also been considered the standard treatment for progressive spinal deformities (e.g., scoliosis) and traumatic injuries. Additionally, lumbar fusion is performed for clearly defined spinal instability. Fusing of the spine is used primarily to eliminate the pain caused by abnormal motion of the vertebrae by immobilizing the faulty vertebrae themselves. Supplementary bone tissue, either from the patient (autograft) or a donor (allograft), is used in conjunction with the body's natural bone growth (osteoblastic) processes, to fuse the vertebrae. There are two main types of lumbar spinal fusion, which may be used in conjunction with each other. Posterolateral fusion places the bone graft between the transverse processes in the back of the spine. These vertebrae are then fixed in place with screws and/or wire through the pedicles of each vertebra, attaching to a metal rod on each side of the vertebrae. Interbody fusion places the bone graft between the vertebrae in the area usually occupied by the intervertebral disc. The fusion then occurs between the endplates of the vertebrae. Using both types of fusion is known as 360-degree fusion. There are three types of interbody fusion: anterior lumbar interbody fusion (ALIF); posterior lumbar interbody fusion (PLIF); and transforaminal lumbar interbody fusion (TLIF). Interbody cages, instrumentation such as plates, pedicle screws, or rods, and osteo-inductive agents such as recombinant human bone morphogenetic protein (rhBMP) may be used to stabilize the spine during the months following surgery, to improve fusion success rates. External factors such as smoking, osteoporosis, certain medications, and heavy activity can prolong or even prevent the fusion process.

The Meyerding Classification Grade of Spondylolisthesis is determined by measuring the degree of slip using standing, neutral lateral radiographs of the lumbar spine. The classification system divides slip into five grades: 0% to 25% is Grade I, 25% to 50% is Grade II, 50% to 75% is Grade III, 75% to 100% is Grade IV, and greater than 100% is Grade V.

Dynamic stabilization, also known as soft stabilization or flexible stabilization, has been proposed as an adjunct or alternative to spinal fusion for the treatment of severe refractory pain due to degenerative spondylolisthesis, or continued severe refractory back pain following prior fusion, sometimes referred to as failed back surgery syndrome. Dynamic stabilization uses flexible materials rather than rigid devices to stabilize the affected spinal segment(s). These flexible materials may be anchored to the vertebrae by synthetic cords or by pedicle screws. Unlike the rigid fixation of spinal fusion, dynamic stabilization is intended to preserve the mobility of the spinal segment.

RATIONALE

Lumbar spinal fusion is a surgical procedure and does not require approval by the FDA. A variety of instrumentation used in lumbar spinal fusion is cleared for marketing by the FDA.

Smoking

Tobacco use is considered a risk factor for poor healing and is associated with non-union. It is well-established that smoking is a preventable cause of morbidity and mortality. The American Academy of Orthopedic Surgeons (AAOS) strongly recommends avoiding use and exposure to tobacco products due to the severe and negative impact on the musculoskeletal system, including the bones, muscle, tendons, and ligaments (AAOS, 2010). Lumbar fusion is in most situations an elective surgery; it is strongly recommended that individuals be in the best physical condition prior to

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 6 of 11

undergoing surgery. A 2011 policy statement published by the International Society of Advancement for Spine Surgery (ISASS) indicated that, while undergoing conservative care prior to surgery, smokers should be encouraged to stop smoking, as smoking aggravates low back pain, is a risk factor for multiple systemic health problems, and increases the risk from poor outcomes of spine surgery. The North American Spine Society (NASS) lists the absence of smoking for at least three months prior to the surgery date in its coverage policy recommendations for lumbar fusion to relieve discogenic low back pain. Anderson et al. (2010) reported that smoking negatively affects fusion mass and results in lower bone mineral density, particularly in the spine. Deyo et al. (2010) evaluated trends and complications in adults who underwent lumbar fusion for spinal stenosis and noted that, not only did major complications increase with increased comorbidity, but there was a substantially greater risk among those with chronic lung disease compared to those without. Particularly with spinal fusion, tobacco use has been associated with increased risk of pseudarthrosis. In addition, tobacco use has been associated with poorer clinical outcomes such as less pain relief, poorer functional rehabilitation, and less overall patient satisfaction (Vogt et al., 2002).

Cotinine, the primary metabolite of nicotine, is currently regarded as the best biomarker of tobacco smoke exposure. Measuring cotinine is preferable to measuring nicotine because cotinine persists longer in the body, with a plasma half-life of about 16 hours. Non-smokers exposed to typical levels of second-hand smoke have serum cotinine levels lower than one ng/ml, with heavy exposure to second-hand smoke producing levels in the 1-10 ng/ml range. Active smokers almost always have levels higher than 10 ng/ml and sometimes higher than 500 ng/ml. Therefore, non-smoking is defined as a serum cotinine level of less than or equal to 10 ng/ml (National Biomonitoring Program, Centers for Disease Control and Prevention, Dec 2013).

Disc herniation/degenerative disc disease (DDD)

Current evidence, which includes a large, multi-center, randomized, controlled trial (RCT) by Weinstein and colleagues known as Spine Patient Outcomes Research Trial (SPORT), supports that surgical treatment with discectomy improves outcomes for lumbar disc herniation with radiculopathy. However, there is no evidence to support that the addition of spinal fusion to discectomy improves outcomes in patients with the sole indication of lumbar disc herniation without instability (e.g., Takeshima et al., 2000, Otani et al., 2014).

W.C. Jacobs and colleagues (2011) conducted a systematic review to assess the effects of surgery versus conservative therapy (including epidural injections) for patients with sciatica due to lumbar disc herniation. RCTs of adults with lumbar radicular pain that evaluated at least one clinically relevant outcome measure (pain, functional status, perceived recovery, lost days of work) were included. In total, five studies were identified, two of which had a low risk of bias. One study compared early surgery with prolonged conservative care followed by surgery, if needed; three studies compared surgery with usual conservative care; and one study compared surgery with epidural injections. Data were not pooled because of clinical heterogeneity and poor reporting of data. One large low-risk-of-bias trial demonstrated that early surgery in patients with six to 12 weeks of radicular pain leads to faster pain relief when compared with prolonged conservative treatment, but there were no differences after one and two years. Another large, low-risk-of-bias trial comparing surgery and usual conservative care found no statistically significant differences on any of the primary outcome measures after one and two years. Future studies should evaluate which patients benefit more from surgery and which from conservative care.

Evidence supporting lumbar fusion as a method of treatment for DDD is limited, and few well-designed clinical studies have supported arthrodesis as superior to non-operative therapy for improving clinical outcomes (e.g., Resnick et al., 2005). When comparing intense rehabilitation and cognitive therapy to lumbar fusion, the reported clinical outcomes demonstrate that lumbar fusion is no more effective than intense rehabilitation combined with cognitive therapy (e.g., Brox et al., 2010; Mirza et al., 2007; Brox et al., 2006; Fairbank et al., 2005). The North American Spine Society (NASS) states that lumbar fusion is not indicated for disc herniation as an adjunct to primary excision of a central or posterolateral disc herniation at any level, in the absence of instability or spondylolisthesis.

Chronic Low Back Pain (CLBP)

A systematic review from 2013 by Saltychev et al. compared lumbar fusion versus conservative treatment in patients with CLBP. The meta-analysis of four trials with a total of 666 patients reported a reduction in the ODI that was -2.91 in favor of lumbar fusion. However, this did not attain statistical significance or the minimal clinically significant difference in

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 7 of 11

ODI of 10 points. The review concluded that there is strong evidence that lumbar fusion does not lead to a clinically significant reduction in perceived disability compared with conservative treatment in patients with CLBP and degenerative spinal disease. The review also concluded that it is unlikely that further research on the subject would alter this conclusion.

T. Ibrahim and colleagues performed a meta-analysis of RCTs to investigate the effectiveness of surgical fusion for the treatment of chronic low back pain compared to non-surgical intervention. The meta-analysis comparison was based on the mean difference in Oswestry Disability Index (ODI) change from baseline to the specified follow-up of patients undergoing surgical versus non-surgical treatment. Of the 58 articles identified, three studies were eligible for primary analysis and one study for sensitivity analysis, with a total of 634 patients. The authors found that surgical fusion for chronic low back pain favored a marginal improvement in the ODI, compared to non-surgical intervention. This difference in ODI was not statistically significant and was of minimal clinical importance. Surgery was found to be associated with a significant risk of complications. Therefore, the cumulative evidence at the present time does not support routine surgical fusion for the treatment of chronic low back pain.

Spinal Stenosis with Spondylolisthesis

The SPORT RCT, reported by Weinstein and Colleagues, compared surgical and nonsurgical treatment for lumbar degenerative spondylolisthesis in two articles dated 2007 and 2009. All patients had neurogenic claudication or radicular leg pain associated with neurologic signs, spinal stenosis shown on cross-sectional imaging, and degenerative spondylolisthesis shown on lateral radiographs, with symptoms persisting for at least 12 weeks. There were 304 patients in a randomized cohort and 303 patients in an observational cohort. About 40% of the randomized cohort crossed over in each direction by two years of follow-up. At the four-year follow-up timepoint, 54% of patients randomized to non-operative care had undergone surgery. Five percent of the surgically-treated patients received decompression only, and 95% underwent decompression with fusion. Analysis was by the treatment that was received, due to the high percentage of crossovers. This analysis, controlled for baseline factors, showed a significant advantage for surgery at up to four years of follow-up for all primary and secondary outcome measures.

Adolescent Idiopathic Scoliosis

Treatment of scoliosis currently depends on three factors: the cause of the condition (idiopathic, congenital, secondary), the severity of the condition (degrees of curve), and the remaining growth expected for the patient at the time of presentation. Children who have vertebral curves measuring between 25° and 40° with at least two years of growth remaining are considered to be at high risk of curve progression. Because severe deformity may lead to compromised respiratory function and is associated with back pain in adulthood, in the U.S., surgical intervention with spinal fusion is typically recommended for curves that progress to 45° or more. (Richards et al., 2005). Long-term follow-up of a large case series by Danielson and Nachemson supports guidelines from the Scoliosis Research Society that fusion can reduce curve progression in patients with curves greater than 40°. This is likely to result in reduced morbidity for treated patients.

Adult Symptomatic Lumbar Scoliosis

No RCTs were identified on the treatment of adult symptomatic lumbar scoliosis with fusion. A cohort study in 2009 by Bridwell et al. reported a prospective, multi-center cohort study that compared operative versus non-operative treatment of adult symptomatic lumbar scoliosis (defined as a minimum Cobb angle of 30°) in 160 consecutively enrolled patients. Operative versus non-operative treatment was decided by the patient and medical team. Non-operative treatment included observation (21%), medications (26%), medications plus physical therapy and/or injections (40%), and other treatment without medications (13%). For analysis, the patients were matched using propensity scores that included baseline Cobb angle, Oswestry Disability Index (ODI), Scoliosis Research Society subscore, and a numerical rating scale for back and leg pain. The percentage of patients who returned for follow-up at two years was higher for operative than non-operative patients (95% vs 45%), though the baseline measures for patients who were lost to follow-up was similar to those who were followed for two years. At the two-year follow-up, non-operative treatment had not improved quality of life or any other outcome measures, while the operative group showed significant improvement in all outcomes.

Isthmic Spondylolisthesis

An RCT compared fusion versus an exercise program for patients with symptomatic isthmic spondylolisthesis. Results of this trial support that the use of fusion for this condition improves functional status, compared with conservative

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 8 of 11

treatment. Moller and Hedlund reported a study of 111 patients with adult isthmic spondylolisthesis who were randomly assigned to posterolateral fusion (with or without instrumentation, n=77) or to an exercise program (n=34). Inclusion criteria for the study were lumbar isthmic spondylolisthesis of any grade, at least one year of low back pain or sciatica, and a severely restricted functional ability. The mean age of patients was 39 years, with a mean age at onset of symptoms of 26 years. At one- and two-year follow-up, functional outcome (assessed by the Disability Rating Index) had improved in the surgery group, but not in the exercise group. Pain scores improved in both groups, but were significantly better in the surgically treated group compared with the exercise group.

Dynamic Stabilization Systems/Devices

There is insufficient research to show that spinal dynamic stabilization devices improve health outcomes for people with disorders of the spine at any level. No clinical guidelines based on research recommend spinal dynamic stabilization devices.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
22533	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); lumbar
22534	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic or lumbar, each additional vertebral segment (List separately in addition to code for primary procedure)
22558	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); lumbar
22585	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); each additional interspace (List separately in addition to code for primary procedure)
22612	Arthrodesis, posterior or posterolateral technique, single level; lumbar (with lateral transverse technique, when performed)
22614	Arthrodesis, posterior or posterolateral technique, single level; each additional vertebral segment (List separately in addition to code for primary procedure)
22630	Arthrodesis, posterior interbody technique, including laminectomy and/or discectomy to prepare interspace (other than for decompression), single interspace; lumbar
22632	Arthrodesis, posterior interbody technique, including laminectomy and/or discectomy to prepare interspace (other than for decompression), single interspace; each additional interspace (List separately in addition to code for primary procedure)
22633	Arthrodesis, combined posterior or posterolateral technique with posterior interbody technique including laminectomy and/or discectomy sufficient to prepare interspace (other than for decompression), single interspace and segment; lumbar

Medical Policy: LUMBAR FUSION FOR ADULTS**Policy Number: 7.01.90****Page: 9 of 11**

Code	Description
22634	Arthrodesis, combined posterior or posterolateral technique with posterior interbody technique including laminectomy and/or discectomy sufficient to prepare interspace (other than for decompression), single interspace and segment; each additional interspace and segment (List separately in addition to code for primary procedure)
22800-22812	Arthrodesis for spinal deformity (code range)
22840-22847	Spinal instrumentation (code range)
22853	Insertion of interbody biomechanical device(s) (e.g., synthetic cage, mesh) with integral anterior instrumentation for device anchoring (e.g., screws, flanges), when performed, to intervertebral disc space in conjunction with interbody arthrodesis, each interspace (List separately in addition to code for primary procedure)
22854	Insertion of intervertebral biomechanical device(s) (e.g., synthetic cage, mesh) with integral anterior instrumentation for device anchoring (e.g., screws, flanges), when performed, to vertebral corpectomy(ies) (vertebral body resection, partial or complete) defect, in conjunction with interbody arthrodesis, each contiguous defect (List separately in addition to code for primary procedure)

*Copyright © 2022 American Medical Association, Chicago, IL***HCPCS Codes**

Code	Description
No codes	

ICD10 Codes

Code	Description
M40.35-M40.37	Flatback syndrome thoracolumbar, lumbar or lumbosacral region (code range)
M41.05-M41.9	Scoliosis (code range: codes ending in 5 are thoracolumbar, ending in 6 are lumbar and ending in 7 are lumbosacral)
M43.00-M43.07	Spondylolysis, thoracolumbar, lumbar or lumbosacral region (code range)
M43.15-M43.17	Spondylolisthesis, thoracolumbar, lumbar or lumbosacral region (code range)
M43.27	Fusion of spine, lumbosacral region
M48.05-M48.07	Spinal stenosis, thoracolumbar, lumbar or lumbosacral region (code range)
M51.06	Intervertebral disc disorders with myelopathy, lumbar region
M53.2X5-M53.2X7	Spinal instabilities, thoracolumbar, lumbar or lumbosacral region (code range)
M53.86-M53.87	Other specified dorsopathies, lumbar or lumbosacral region (code range)
M96.0	Pseudarthrosis after fusion or arthrodesis
M96.1	Postlaminectomy syndrome, not elsewhere classified
S32.000A-S32.059S	Lumbar fracture (code range)

REFERENCES

Abdu RW, et al. Reoperation for Recurrent Intervertebral Disc Herniation in the Spine Patient Outcomes Research Trial. *Spine* July 2017; 42(14) 1106-1114.

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 10 of 11

*Anderson T, et al. Smoking as a predictor of negative outcome in lumbar spinal fusion. Spine 2001 Dec 1;26(23):2623-8.

Blue Cross and Blue Shield Association. Lumbar Spinal Fusion. Medical Policy Reference Manual Policy #7.01.141. 2021 Nov 01.

*Bridwell KH, et al. Does treatment (nonoperative and operative) improve the two-year quality of life in patients with adult symptomatic lumbar scoliosis: a prospective multicenter evidence-based medicine study. Spine (Phila Pa 1976) Sep 15 2009;34(20):2171-2178.

*Brox JJ, et al. Lumbar instrumented fusion compared with cognitive intervention and exercises in patients with chronic back pain after previous surgery for disc herniation: a prospective randomized controlled study. Pain 2006 May;122(1-2):145-55.

*Carreon LY, et al. Fusion and nonsurgical treatment for symptomatic lumbar degenerative disease: a systematic review of Oswestry Disability Index and MOS Short Form-36 outcomes. Spine J 2008 Sep-Oct;8(5):747-55.

*Coe M, et al. The role of fusion for discogenic axial back pain without associated leg pain, spondylolisthesis or stenosis: an evidence-based review. Semin Spine Surg 2009 Dec;21(4):246-56.

Diniz JM and Botelho RV. Is fusion necessary for thoracolumbar burst fracture treated with spinal fixation? A systematic review and meta-analysis. J Neurosurg Spine 2017 Nov;27(5):584-592.

Donnelly CJ, et al. Current incidence of adjacent segment pathology following lumbar fusion versus motion-preserving procedures: a systematic review and meta-analysis of recent projections. The Spine Journal 2020 Oct; 20(10): 1554-1565.

Forst P, et al. A randomized, controlled trial of fusion surgery for lumbar stenosis. New Eng J Med 2016 April 14;374(15):1413-1423.

Ghogawala Z, et al. Laminectomy plus fusion versus laminectomy alone for lumbar spondylolisthesis. New Eng J Med 2016 April 14;374(15):1424-1434.

Giang G, et al. Evaluating outcomes of stand-alone anterior lumbar interbody fusion: A systematic review. World Neurosurg 2017 Aug;104:259-271.

Gaudin D, et al. Considerations in spinal fusion surgery for chronic lumbar pain: Psychosocial factors, rating scales, and perioperative patient education-A review of the literature. World Neurosurg 2017 Feb;98:21-27.

*Institute for Clinical Systems Improvement. Healthcare guideline. Adult acute and subacute low back pain. 2018 Mar [<https://www.icsi.org/guideline/low-back-pain/>] accessed 1/13/22.

International Society for the Advancement of Spine Surgery (ISASS). Policy statement for lumbar spinal fusion surgery. 2011 Jul 15 [https://www.isass.org/pdf/2011-07-15_policy_statement_lumbar_surgery.pdf] accessed 1/13/22.

Jackson KL 2nd, et al. The Effects of Smoking and Smoking Cessation on Spine Surgery: A Systematic Review of the Literature. Global Spine J 2016 Nov;6(7):695-701.

Jackson KL 2nd and Devine JG. The effects of obesity on spine surgery: A systematic review of the literature. Global Spine J 2016 Jun;6(4):394-400. doi: 10.1055/s-0035-1570750.

Kalakoti P, et al. Impact of psychiatric comorbidities on short-term outcomes following intervention for lumbar degenerative disc disease. Spine (Phila Pa 1976) 2018 Oct 1;43(19):1363-1371.

*Kreiner DS, et al. An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spinal stenosis (update). Spine J 2013;13(7):734-43.

McCunniff PT, et al. Smoking is associated with increased blood loss and transfusion use after lumbar spinal surgery. Clin Orthop Relat Res 2016 April;474(4):1019-1025.

McGirt MJ, et al. An analysis from the Quality Outcomes Database, Part 1. Disability, quality of life, and pain outcomes following lumbar spine surgery: predicting likely individual patient outcomes for shared decision-making. J Neurosurg Spine 2017 Oct;27(4):357-369.

Medical Policy: LUMBAR FUSION FOR ADULTS

Policy Number: 7.01.90

Page: 11 of 11

*Moller H and Hedlund R. Surgery versus conservative management in adult isthmic spondylolisthesis--a prospective randomized study: part 1. Spine (Phila Pa 1976) Jul 1 2000;25(13):1711-1715.

Moreau S, et al. Reduction and fusion in high-grade L5-S1 spondylolisthesis by a single posterior approach. Results in 50 patients. Orthop Traumatol Surg Res 2016 April;102(2):233-237.

North American Spine Society. NASS coverage policy recommendations: Lumbar fusion. 2014.

North American Spine Society. Diagnosis and Treatment of Lumbar Disc Herniation with Radiculopathy: Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care. 2012.

[<https://www.spine.org/Documents/ResearchClinicalCare/Guidelines/LumbarDiscHerniation.pdf>] accessed 1/13/22.

North American Spine Society. Clinical practice guidelines. Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis (Revised 2011).

[<https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/LumbarStenosis.pdf>] accessed 1/13/22.

North American Spine Society. Clinical practice guidelines. Diagnosis and Treatment of Degenerative Lumbar Spondylolisthesis. 2014 Revised

[<https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/Spondylolisthesis.pdf>] accessed 1/13/22.

O'Donnell JA, et al. Treatment of recurrent lumbar disc herniation with or without fusion in workers' compensation subjects. Spine (Phila Pa 1976) 2017 Jul 15;42(14):E864-E870.

Pereira BJ, et al. Spinal surgery for degenerative lumbar spine disease: predictors of outcome. Clin Neurol Neurosurg 2016 Jan;140:1-5.

*Richards BS, et al. Standardization of criteria for adolescent idiopathic scoliosis brace studies: SRS Committee on Bracing and Nonoperative Management. Spine (Phila Pa 1976) Sep 15 2005;30(18):2068- 2075; discussion 2076-2067.

*Saltychev M, et al. Lumbar fusion compared with conservative treatment in patients with chronic low back pain: a meta-analysis. Int J Rehabil Res Mar 2014;37(1):2-8.

*Samuel S, et al. Fusion versus conservative management for low-grade isthmic spondylolisthesis. Cochrane Database System Rev 2012;10:CD010150.

Takaoka H, et al. Comparison between intervertebral oblique lumbar interbody fusion and transforaminal lumbar interbody fusion: a multicenter study. Scientific Reports 2021;11: 1-10.

Wang G, et al. Evaluation and surgical management of adult degenerative scoliosis associated with lumbar stenosis. Medicine 2016 April;95(15):e3394.

*Weinstein JN, et al. Surgical versus nonoperative treatment for lumbar disc herniation: four year results for the Spine Patient Outcomes Research Trial (SPORT). Spine 2008 Dec 1;33(25):2789.

Zhang C, et al. Adjacent segment degeneration versus disease after lumbar spine fusion for degenerative pathology: a systematic review with meta-analysis of the literature. Clin Spine Surg 2016 Feb;29(1):21-29

*Key Article

KEY WORDS

Degenerative disc disease, Disc herniation, Lumbar arthrodesis, Lumbar fusion, Spinal stenosis, spondylodesis, spondylosyndesis, Spondylolisthesis

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based upon our review, lumbar fusion is not addressed in National or regional CMS coverage determinations or policies.