

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION
Policy Number	7.01.51
Category	Technology Assessment
Effective Date	11/15/01
Revised Date	09/19/02, 09/18/03, 07/15/04, 07/21/05, 05/18/06, 04/19/07, 06/19/08, 05/28/09, 04/22/10, 03/17/11, 03/15/12, 06/19/14, 08/20/15, 10/20/16, 10/19/17, 06/21/18, 12/20/18
Archived Date	(03/21/13-06/19/14)
Product Disclaimer	<ul style="list-style-type: none"> • <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i> • <i>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</i> • <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i>

POLICY STATEMENT

- I. Based upon our criteria and assessment of peer-reviewed literature, implantation of a dorsal column spinal cord stimulator (i.e. low frequency or high frequency) has been medically proven to be effective and therefore, **medically appropriate** for treatment of patients with failed back syndrome (FBS) with intractable neuropathic leg pain for the following:
- A. A short term trial (e.g., greater than 48 hours) of spinal cord stimulation (i.e. low frequency or high frequency (HF 10 SCS) is considered medically necessary for the treatment of chronic intractable pain secondary to failed back surgery syndrome (FBSS) with intractable neuropathic leg pain when ALL of the following criteria are met:
 1. Failure of at least six consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive therapy, and activity lifestyle modification);
 2. Surgical intervention is not indicated or patients do not wish to proceed with spinal surgery; and
 3. An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate placement of the device.
 - B. Permanent implantation of a spinal cord stimulator (i.e. low frequency or high frequency (HF 10 SCS) is considered medically necessary for the treatment of chronic intractable pain secondary to FBSS with intractable neuropathic leg pain when at least a 50% reduction in pain has been demonstrated during a short-term trial of SCS.
- II. Based upon our criteria and assessment of peer-reviewed literature, use of a low frequency dorsal column spinal cord stimulator has been medically proven to be effective and therefore, **medically appropriate** for treatment of patients with Complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) for the following:
- A. A short term trial (e.g., greater than 48 hours) of a low frequency dorsal column spinal cord stimulation of a low frequency spinal cord stimulator (SCS) is considered medically necessary for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) when all of the following criteria are met:
 1. Failure of at least six consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive therapy, and activity lifestyle modification);
 2. Surgical intervention is not indicated or patients do not wish to proceed with spinal surgery;
 3. An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would negatively impact the success of a SCS or contraindicate its placement.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 2 of 13

- B. Permanent implantation of a low frequency dorsal column spinal cord stimulator is considered medically necessary for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) when at least a 50% reduction in pain has been demonstrated during a short-term trial of SCS
- III. Based upon our criteria and assessment of peer-reviewed literature, use of a low frequency dorsal column spinal cord stimulator has been medically proven to be effective and therefore, **medically appropriate** for treatment of patients with chronic critical limb ischemia (CLI) for the following:
- A. A short term trial (e.g., greater than 48 hours) of a low frequency dorsal column spinal cord stimulation of a low frequency spinal cord stimulator (SCS) is considered medically necessary for the treatment of chronic, intractable pain secondary to chronic critical limb ischemia (CLI) when both of the following criteria are met
1. Failure of available conventional multidisciplinary medical (e.g., pharmacological, physical therapy) and surgical management (e.g., revascularization); and
 2. An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would negatively impact the success of a SCS or contraindicate its placement.
- B. Permanent implantation of a low frequency dorsal column spinal cord stimulator is considered medically necessary for the treatment of chronic, intractable pain secondary to chronic critical limb ischemia (CLI) when a beneficial clinical response from a temporarily implanted electrode has been demonstrated prior to consideration of permanent implantation.
- IV. Based upon our criteria and assessment of peer-reviewed literature, use of a low frequency dorsal column spinal cord stimulator has been medically proven to be effective and therefore, **medically appropriate** for treatment of patients with chronic, intractable pain secondary to chronic stable angina pectoris for myocardial ischemia for the following:
- A. A short term trial (e.g., greater than 48 hours) of a low frequency dorsal column spinal cord stimulation of a low frequency spinal cord stimulator (SCS) is considered medically necessary for the treatment of chronic, intractable pain secondary to chronic stable angina pectoris/myocardial ischemia when all of the following criteria are met:
1. Angina pectoris is Canadian Cardiovascular Society (CCS) functional class III or class IV;
 2. An attestation from the Individual's treating cardiologist confirming that the individual has coronary artery disease (CAD) and is not a suitable candidate for a revascularization procedure;
 3. Optimal pharmacological treatment using anti-anginal medications (e.g., long-acting nitrates, beta-adrenergic blockers, or calcium-channel antagonists) has failed to adequately improve anginal symptoms; and
 4. An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate its placement of the device.
- B. Permanent implantation of a low frequency dorsal column spinal cord stimulation of a low frequency spinal cord stimulator (SCS) is considered medically necessary for the treatment of chronic, intractable pain secondary to chronic stable angina pectoris for myocardial ischemia when a beneficial clinical response from a temporarily implanted electrode has been demonstrated prior to consideration of permanent implantation.
- V. Based upon our criteria and assessment of peer-reviewed literature, the replacement of an existing high frequency or low frequency dorsal column spinal cord stimulator (SCS) and/or battery/generator is considered **medically necessary** for an individual when the existing stimulator and/or battery/generator is malfunctioning, cannot be repaired, and is no longer under warranty.
- VI. Replacement of a functioning low frequency dorsal column spinal cord stimulator with a high frequency spinal cord stimulator is considered **not medically necessary**.
- VII. Based upon our criteria and assessment of peer-reviewed literature, implantation of a low frequency dorsal column spinal cord stimulator has not been medically proven to be effective and is considered **investigational**, for all other indications, including but not limited to:
- A. Post-amputation pain (phantomalimb pain)

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 3 of 13

- B. Post-herpetic neuralgia
- C. Peripheral neuropathy
- D. Dysesthesias involving the lower extremities secondary to spinal cord injury.

VIII. Based upon our criteria and assessment of peer-reviewed literature, implantation of a high frequency spinal cord stimulator has not been medically proven to be effective and is considered **investigational** for all other indications.

IX. Based upon our criteria and assessment of peer-reviewed literature, peripheral nerve stimulation, **dorsal root ganglion** stimulation, including peripheral nerve field stimulation, is considered **investigational** for treatment of acute or chronic pain conditions, including treatment of the following;

- A. Failed back surgery syndrome (FBSS) with intractable neuropathic leg pain
- B. Complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD)
- C. Chronic Critical Limb Ischemia (CLI)
- D. Chronic Stable Angina Pectoris
- E. Post-amputation pain (phantom limb pain)
- F. Post-herpetic neuralgia
- G. Peripheral neuropathy; and
- H. Dysesthesias involving the lower extremities secondary to spinal cord injury

Refer to Corporate Medical Policy #11.01.03 regarding Experimental and Investigational Services.

*This medical policy does not address **occipital nerve stimulation** for chronic migraines or occipital neuralgia. In occipital nerve stimulation the neurostimulator delivers electrical impulses via insulated lead wires tunneled under the skin near the occipital nerves at the base of the head. Currently, there is no FDA approved device for this indication.*

POLICY GUIDELINES

- I. The implantation of a spinal cord stimulator is used only as a last resort. Other treatment modalities (pharmacological, surgical, psychological, or physical, if applicable) need to have been tried and failed or have been judged unsuitable or contraindicated. Duration of refractory pain is six months or greater.
- II. Documentation must reflect an objective measure of a 50% reduction in pain scores with a temporarily implanted electrode in order to precede permanent implantation.
- III. Patients are to be carefully screened, evaluated, and diagnosed by a multidisciplinary team prior to application of these therapies. This evaluation may include a psychological evaluation to exclude any major mental disability or drug habituation that would negatively influence the outcome of the treatment. *Please to Refer to Corporate Medical Policy #3.01.02 regarding Psychological Testing.*
- IV. The Federal Employees Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity

DESCRIPTION

Spinal cord stimulation (SCS) is used to treat chronic back and extremity pain and consists of electrical stimulation of the dorsal columns by electrodes implanted in the epidural space. The neurophysiology of pain relief after spinal cord stimulation is uncertain, but may be related to either activation of an inhibitory system or blockage of facilitatory circuits. Spinal cord stimulation devices consist of implantable electrodes, a receiver/transducer and a programmable transmitter that may be worn externally or implanted. Implantation of the spinal cord stimulator is typically a two-step process. Initially the electrode(s) is temporarily implanted in the epidural space, allowing a trial period of stimulation. This trial

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 4 of 13

period will typically last for a period of 3 to 7 days. Once treatment effectiveness has been established, the electrode(s) and receiver/transducer are permanently implanted. Successful spinal cord stimulation may require extensive programming to determine the optimum levels of stimulation to provide pain relief. There are two basic types of power source. In 1 type, the power source (battery) can be surgically implanted. In another, a radio-frequency receiver is implanted and the power source is worn externally with an antenna over the receiver. Totally implantable systems are most commonly used.

Spinal cord stimulation has been utilized in a variety of refractory neuropathic pain conditions, including pain associated with failed back syndrome, arachnoiditis, peripheral neuropathy and complex regional pain syndrome. Complex regional pain syndrome (CRPS) is a chronic pain condition most often affecting one of the limbs (arms, legs, hands, or feet), usually after an injury or trauma to that limb. CRPS is believed to be caused by damage to, or malfunction of, the peripheral and central nervous systems. The central nervous system is composed of the brain and spinal cord, and the peripheral nervous system involves nerve signaling from the brain and spinal cord to the rest of the body. CRPS is characterized by prolonged or excessive pain and mild or dramatic changes in skin color, temperature, and/or swelling in the affected area. There are two similar forms, called CRPS-I and CRPS-II, with the same symptoms and treatments. CRPS-II (previously called causalgia) is the term used for patients with confirmed nerve injuries. Individuals without confirmed nerve injury are classified as having CRPS-I (previously called reflex sympathetic dystrophy syndrome). People with CRPS also experience constant or intermittent changes in temperature, skin color, and swelling of the affected limb. This is due to abnormal microcirculation caused by damage to the nerves controlling blood flow and temperature. An affected arm or leg may feel warmer or cooler compared to the opposite limb. The skin on the affected limb may change color, becoming blotchy, blue, purple, pale, or red.

Spinal cord stimulation is generally not effective in treating nociceptive pain (pain resulting from irritation, as opposed to damage to the nerves) and central deafferentation pain (pain related to central nervous system damage from a stroke or spinal cord injury).

It is recommended that candidates for SCS undergo a psychological evaluation prior to surgery. The purpose of the evaluation is to assess the potential role that psychological factors (e.g., anxiety, depression, underlying mental illness) may have in influencing the success of surgery and to offer appropriate recommendations with regard to psychological management.

Spinal cord stimulation has also been investigated as a treatment for pain associated with cervical trauma or disc herniation, chronic refractory angina pectoris and critical limb ischemia in patients who are not candidates for revascularization procedures.

High frequency spinal cord stimulation, also referred to as kilohertz frequency spinal cord stimulation or HF10, is a type of spinal cord stimulation (SCS) providing a higher frequency than traditional spinal cord stimulator systems. The HF10 SCS uses low amplitude, high frequency, and short duration pulses. HF10 SCS does not generate paresthesia and operates at a frequency of 10,000 Hz to provide pain relief in comparison to traditional SCS systems, which operate at a frequency in the range of 40-60 Hz and do generate paresthesia. As an alternative to traditional dorsal spinal column stimulation, HF 10 SCS is proven safe and effective for treatment of chronic, intractable low back and leg pain in patients with failed back surgery syndrome.

Dorsal root ganglion stimulation or peripheral nerve stimulation involves implantation of electrodes near or on a peripheral nerve to reduce pain. Peripheral nerve field stimulation is a technology that involves placement of electrodes subcutaneously within an area of maximal pain, with the objective of stimulating a region of affected nerves to reduce pain. Depending on the targeted nerve, leads may be placed percutaneously just under the skin or via an open approach for larger deeper peripheral nerves. Similar to SCS, a short term trial is required prior to permanent implantation of a

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 5 of 13

generator. The use of these technologies, used alone or in combination with SCS for the treatment of pain conditions is under investigation

RATIONALE

Traditional stimulation

Totally implantable spinal cord (dorsal column) stimulator systems are regulated by the FDA as class III pre-market-approval (PMA) devices. Several devices have received FDA PMA approval. Examples of these devices include, but are not limited to, the Precision™ Spinal Cord Stimulator System, and the Genesis™ IPG System. Systems with external transmitters are regulated by the FDA as Class II 510(K) devices. The FDA gave 510 K approval for Advanced Neuromodulation systems to market their Renew spinal cord stimulator, to Medtronic for its Spinal Cord and Peripheral Nerve Stimulation Systems, X-trel®3 and Synergy®; Spinal Cord Stimulation Systems, and to Micronet Medical, Inc for its Axxess Spinal Cord Stimulation Lead. St. Jude Medical has also received FDA approval for its Protege MRI™ spinal cord stimulation system.

There is sufficient evidence in the peer -reviewed literature to permit conclusions that the technology provides significant and sustained relief of pain with minimal side effects in appropriately selected patients with chronic nonmalignant pain. Studies investigating the effectiveness of SCS as a treatment for patients with chronic back/extremity pain report successful management of pain, a substantial decrease in narcotic use and an improvement in the quality of life. Studies support the use of spinal cord stimulation for patients with CRPS in the upper extremities through outcomes that demonstrate reduction in pain intensity and increased quality of life (e.g., Harke, et al. 2005; Kemler, et al. 2006; Kumar, et al. 2011; Geurts, et al. 2013).

One essential step toward the effective use of SCS in potential patients is a trial of the system through percutaneous lead placement. This trial will determine the effectiveness in relieving pain (greater than 50% pain relief) and improving the quality of life in patients with refractory neuropathic pain.

Literature exists to support the value of a presurgical psychological evaluation to identify factors that may adversely impact functional outcomes after spinal cord stimulation (DM Doleys 2006, B Van Dorsten 2006, DR Heckler, et al. 2007, J Celestin, et al. 2009).

There is evidence to favor spinal cord stimulation over standard conservative treatment to improve limb salvage and clinical situation in patients with inoperable chronic critical leg ischemia.

Studies found that spinal cord stimulation improved both the quality of life and cardiac parameters of patients with refractory angina pectoris.

High-frequency stimulation

Nevro (Menlo Park, Calif) has gained FDA 510(k) clearance for its Senza spinal cord stimulation system intended for chronic pain treatment in May 2015. The device administers the company's HF10 therapy in the trunk and/or limbs, which treats unilateral or bilateral pain related to failed back surgery, intractable low back pain and leg pain. The therapy is the only SCS therapy FDA-indicated to alleviate pain without paresthesia (a constant tingling sensation associated with traditional SCS techniques)

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION**Policy Number: 7.01.51****Page: 6 of 13****CPT Codes**

Code	Description
63650	Percutaneous implantation of neurostimulator electrode array; epidural
63655	Laminectomy for implantation neurostimulator electrode plate/paddle; epidural
63661	Removal of spinal neurostimulator electrode percutaneous array(s), including fluoroscopy, when performed
63662	Removal of spinal neurostimulator electrode plate/paddle(s) placed via laminotomy or laminectomy, including fluoroscopy when performed
63663	Revision including replacement, when performed, of spinal neurostimulator electrode percutaneous array(s) including fluoroscopy, when performed
63664	Revision including replacement, when performed, of spinal neurostimulator electrode plate/paddle(s) placed via laminotomy or laminectomy, including fluoroscopy, when performed
63685	Insertion or replacement of spinal neurostimulator pulse generator or receiver, direct or inductive coupling
63688	Revision or removal of implanted spinal neurostimulator pulse generator or receiver
95970-95973	Neurostimulator programming and analysis (code range)

*Copyright © 2019 American Medical Association, Chicago, IL***HCPCS Codes**

Code	Description
C1820	Generator, neurostimulator (implantable), non-high frequency with rechargeable battery and charging system
C1822	Generator, neurostimulator (implantable), high frequency with rechargeable battery and charging system
L8679	Implantable neurostimulator pulse generator, any type
L8680	Implantable neurostimulator electrode, each
L8681	Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only
L8682	Implantable neurostimulator radiofrequency receiver
L8683	Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
L8685	Implantable neurostimulator pulse generator, single array, rechargeable, includes extension
L8686	Implantable neurostimulator pulse generator, single array, non-rechargeable, includes extension
L8687	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
L8688	Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension
L8689	External recharging system for battery (internal) for use with implantable neurostimulator, replacement only
L8695	External recharging system for battery (external) for use with implantable neurostimulator, replacement only

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 7 of 13

ICD10 Codes

Code	Description
Multiple diagnosis codes	

REFERENCES

Abu Dabrh AM, et al. Nonvascularization-based treatments in patients with severe or critical limb ischemia. J Vasc Surg 2015 Nov;62(5):1330-1339.

Al-Kaisy A, et al. Sustained effectiveness of 10kHz high-frequency spinal cord stimulation for patients with chronic, low back pain: 24-month results of a prospective multicenter study. Pain Med 2014 Mar;15(3):347-54.

Al-Kaisy A, et al. 10 kHz high-frequency spinal cord stimulation for chronic axial low back pain in patients with no history of spinal surgery: a preliminary, prospective, open-label and proof-of-concept study. Neuromodulation 2017 Jan;20(1):63-70.

*Amann W, et al. Spinal cord stimulation in the treatment of nonreconstructable stable critical leg ischemia: results of the European Peripheral Vascular Disease Outcome Study (SCS-EPOS). Eur J Vasc Endovasc Surg 2003 Sep;26(3):280-6.

*American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management. Anesthesiol 2010 Apr;112(4):1-24.

American Society of Interventional Pain Physicians. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. 2009 [<http://www.painphysicianjournal.com/> 2009/july/2009;12;699-802.pdf] accessed 7/18/17.

*Andrell P, et al. Long-term effects of spinal cord stimulation on angina symptoms and quality of life in patients with refractory angina pectoris- results from the European Angina Registry Link Study (EARL). Heart 2010 Jul;96(14):1132-6.

Arcioni R, et al. Cervical 10 kHz spinal cord stimulation in the management of chronic, medically refractory migraine: A prospective, open-label, exploratory study. Eur J Pain 2016 Jan;20(1):70-78.

*Bala MM, et al. Systematic review of the (cost-) effectiveness of spinal cord stimulation for people with failed back surgery syndrome. Clin J Pain 2008 Nov-Dec;24(9):741-56.

Bicket MC, et al. High-frequency spinal cord stimulation for chronic pain: Pre-clinical overview and systematic review of controlled trials. Pain Med 2016; 17(12): 2326-36.

BlueCross BlueShield Association. Spinal cord stimulation. Medical Policy Reference Manual Policy # 7.01.25. 2017 July 13.

*Bondesson S, et al. Comparison of patients undergoing enhanced external counterpulsation and spinal cord stimulation for refractory angina pectoris. Coron Artery Dis 2008 Dec;19(8):627-34.

*Borjesson M, et al. Spinal cord stimulation in severe angina pectoris- a systematic review based on the Swedish Council on technology assessment in health care report on long-standing pain. Pain 2008 Dec;140(3):501-8.

*British Pain Society. Consensus statement. Spinal cord stimulation for the management of pain: Recommendations for best clinical practice. 2005 Mar [http://www.britishpainsociety.org/pdf/scs_2005.pdf] accessed 7/18/17.

*Buchser E, et al. Spinal cord stimulation for the management of refractory angina pectoris. J Pain Symptom Manag 2006 Apr;31(4 Suppl):S36-42.

Buffenoir K, et al. Spinal cord stimulation of the conus medullaris for refractory pudendal neuralgia: a prospective study of 27 consecutive cases. Neurourol Urodyn 2015 Feb;34(2):177-182.

Campbell CM, et al. Psychological screening/phenotyping as predictors for spinal cord stimulation. Curr Pain Headache Rep 2013 Jan;17(1):307.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 8 of 13

- *Celestin J, et al. Pretreatment psychosocial variables as predictors of outcomes following lumbar surgery and spinal cord stimulation: a systematic review and literature synthesis. Pain Med 2009 May-Jun;10(4):639-53.
- *Deer TR. Current and future trends in spinal cord stimulation for chronic pain. Cur Pain Headach Rep 2001;5:503-9.
- *Deer TR, et al. Spinal cord stimulation for refractory angina pectoris and peripheral vascular disease. Pain Physician 2006 Oct;9(4):347-52.
- Deer TR, et al. A prospective study of dorsal root ganglion stimulation for relief of chronic pain. Neuromodulation 2013 Jan-Feb;16(1):67-71.
- Deer TR, et al. Effectiveness of cervical spinal cord stimulation for the management of chronic pain. Neuromodulation 2014 Apr;17(3):265-71.
- Deer TR, et al. The appropriate use of neurostimulation of the spinal cord and peripheral nervous system for the treatment of chronic pain and ischemic diseases: the Neuromodulation Appropriateness Consensus Committee. Neuromodulation 2014 Aug;17(6):515-50.
- Deer TR, et al. Dorsal root ganglion stimulation yielded higher treatment success rate for complex regional pain syndrome and causalgia at 3 and 12 months: a randomized comparative trial. Pain April 2017;158(4):669-681.
- De Andres J, et al. Prospective, Randomized Blind Effect-on-Outcome Study of Conventional vs High-Frequency Spinal Cord Stimulation in Patients with Pain and Disability Due to Failed Back Surgery Syndrome. Pain Medicine 2017;18: 2401-2421.
- De Ridder D, et al. Burst spinal cord stimulation for limb and back pain. World Neurosurg 2013 Nov;80(5):642-649.
- deVos CC, et al. Burst spinal cord stimulation evaluated in patients with failed back pain surgery syndrome and painful diabetic neuropathy. Neuromodulation 2014 Feb;17(2):152-9.
- *Di Pede F, et al. Immediate and long-term clinical outcome after spinal cord stimulation for refractory stable angina pectoris. Am J Cardiol 2003 Apr 15;91(8):951-5.
- *Doleys DM. Psychological factors in spinal cord stimulation therapy: brief review and discussion. Neurosurg Focus 2006 Dec 15;21(6):E1.
- Dworkin RH, et al. Interventional management of neuropathic pain: NeuPSIG recommendation. Pain 2013 Nov;154(11):2249-61.
- *Eddicks S, et al. Thoracic spinal cord stimulation improves functional status and relieves symptoms in patients with refractory angina pectoris: the first placebo-controlled study. Heart 2007 May;93(5):585-90.
- *Ekre O, et al. Long-term effects of spinal cord stimulation and coronary artery bypass grafting on quality of life and survival in the ESBY study. Eur Heart J 2002 Dec;23(24):1938-45.
- Eldabe S, et al. The effectiveness and cost-effectiveness of spinal cord stimulation for refractory angina (RASCAL Study): a pilot randomized controlled trial. Neuromodulation 2016 Jan;19(1):60-70.
- Food and Drug Administration. Stimwave Technologies Freedom Spinal Cord Stimulator (SCS System). [http://www.accessdata.fda.gov/cdrh_docs/pdf14/K141399.pdf] accessed 7/18/17.
- *Frey ME, et al. Spinal cord stimulation for patients with failed back syndrome: A systematic review. Pain Physician 2009 Mar-Apr;12(2):379-97.
- Geurts JW, et al. Spinal cord stimulation for complex regional pain syndrome type I: a prospective cohort study with long-term follow-up. Neuromodulation 2013 Nov-Dec;16(6):523-9.
- *Gibbons RJ, et al. ACC/AHA 2002 Guideline update for the management of patients with chronic stable angina-summary article. Circ 2003 Jan;107:149-58.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 9 of 13

Grider JS, et al. Effectiveness of spinal cord stimulation in chronic spinal pain: a systematic review. Pain Physician 2016 Jan;19(1):E33-E54.

*Harke H, et al. Spinal cord stimulation in sympathetically maintained complex regional pain syndrome type I with severe disability. A prospective clinical study. Eur J Pain 2005 Aug;9(4):363-73.

*Harney D, et al. Complex regional pain syndrome: the case for spinal cord stimulation (a brief review). Injury 2005 Mar;36(3):357-62.

*Heckler DR, et al. Presurgical Behavioral Medicine Evaluation (PBME) for implantable devices for pain management: a 1-year prospective study. Pain Pract 2007 Jun;7(2):110-22.

*Henning H, et al. Spinal cord stimulation in postherpetic neuralgia and in acute herpes zoster pain. Anes Analges 2002 Mar;94(3):694-700.

Hou S, et al. A systematic evaluation of burst spinal cord stimulation for chronic back and leg pain. Neuromodulation 2016 June;19(4):398-405.

Institute for Clinical Systems Improvement (ICSI). Assessment and management of chronic pain. 2009 Nov [http://www.icsi.org/guidelines_and_more/gl_os_prot/musculo-skeletal/pain__chronic__assessment_and_management_of_14399/pain__chronic__assessment_and_management_of_14400.html] accessed 7/18/17.

Kapural L, et al. Novel 10-kHz high frequency therapy (HF10 Therapy) is superior to traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: The SENZA-RCT randomized controlled trial. Anesthesiology 2015;123(4):851-860.

Kapural L, et al. Comparison of 10-kHz high-frequency and traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: 24-month results from a multicenter, randomized, controlled pivotal trial. Neurosurgery 2016 Nov;79(5):667-677.

Kelly GA, et al. The impact of spinal cord stimulation on physical function and sleep quality in individuals with failed back surgery syndrome: a systematic review. Eur J Pain 2012 Jul;16(6):793-802.

*Kemler MA, et al. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. NEJM 2000 Aug 31;343(9):618-24.

*Kemler MA, et al. Impact of spinal cord stimulation on sensory characteristics in complex regional pain syndrome type I: a randomized trial. Anesthesiol 2001 Jul;95(1):72-80.

*Kemler MA, et al. Spinal cord stimulation for chronic reflex sympathetic dystrophy- five-year follow-up. N Engl J Med 2006 Jun 1;354(22):2394-6.

*Klomp HM, et al. Spinal cord stimulation is not cost-effective for non-surgical management of critical limb ischaemia. Eur J Vasc Endovas Surg 2006 May;31(5):500-8.

*Klomp HM, et al. What is the evidence on efficacy of spinal cord stimulation in (subgroups of) patients with critical limb ischemia? Ann Vasc Surg 2009 May-Jun;23(3):355-63.

Knife TM, et al. High frequency (10 kHz) or burst stimulation in failed back surgery syndrome patients with predominant back pain: preliminary data from a prospective observational study. Neuromodulation 2016 April;19(3):268-275.

Kriek N, et al. Comparison of tonic spinal cord stimulation, high-frequency and burst stimulation in patients with complex regional pain syndrome: a double-blind, randomized placebo controlled trial. BMC Musculoskelet Disord 2015 Aug 25;16:222.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 10 of 13

*Kumar K, et al. Spinal cord stimulation in treatment of chronic benign pain: challenges in treatment planning and present status, a 22-year experience. Neurosurg 2006 Mar;58(3):481-96.

*Kumar K, et al. Spinal cord stimulation versus conventional medical management for neuropathic pain: a multicentre randomised controlled trial in patients with failed back surgery syndrome. Pain 2007 Nov;132(1-2):179-88.

*Kumar K, et al. The effects of spinal cord stimulation in neuropathic pain are sustained: a 24-month follow-up of the prospective randomized controlled multicenter trial of the effectiveness of spinal cord stimulation. Neurosurg 2008 Oct;63(4):762-70.

*Kumar K, et al. Spinal cord stimulation is effective in management of complex regional pain syndrome I: fact or fiction. Neurosurgery 2011 Sep;69(3):566-78.

Lad SP, et al. Utilization of spinal cord stimulation in patients with failed back surgery syndrome. Spine 2014 May 20;39(12):E719-27.

Lambro G, et al. Safety and efficacy of cervical 10 kHz spinal cord stimulation in chronic refractory primary headaches.; a retrospective case series. J headache Pain 2016 Dec;17(1):66.

*Lanza GA, et al. Effect of spinal cord stimulation on spontaneous and stress reduced angina and ischemic-like ST-segment depression in patients with cardiac syndrome X. Eur Heart J 2005 Jan 9.

*Lanza GA, et al. Spinal cord stimulation for the treatment of refractory angina pectoris: a multicenter randomized single-blind study (the SCS-ITA trial). Pain 2011 Jan;152(1):45-52.

Liem L, et al. A multicenter, prospective trial to assess the safety and performance of the spinal modulation dorsal root ganglion neurostimulator system in the treatment of chronic pain. Neuromodulation 2013 Sept-Oct;16(5):471-482.

Liem L, et al. One year outcomes of spinal cord stimulation of the dorsal root ganglion in the treatment of chronic neuropathic pain. Neuromodulation 2015 Jan;18(1):41-48.

Lihua P, et al. Spinal cord stimulation for cancer-related pain in adults. Cochrane Database Syst Rev 2013 Feb 28;1:CD009389.

*Mailis-Gagnon A, et al. Spinal cord stimulation for chronic pain. Cochrane Database Syst Review 2004;(3): CD003783.

*Manca A, et al. Quality of life, resource consumption and costs of spinal cord stimulation versus conventional medical management in neuropathic pain patients with failed back surgery syndrome (PROCESS trial). Eur J Pain 2008 Nov;12(8):1047-58.

Mailis A, et al. Evidence-based guideline for neuropathic pain interventional treatments: spinal cord stimulation, intravenous infusions, epidural injections and nerve blocks. Pain Res Manag 2012 May-Jun;17(3):150-8.

Manchikani L, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part I: introduction and general considerations. Pain Physician 2013 Apr;16(2 Suppl):S1-48.

Manchikani L, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. Pain Physician 2013 Apr;16(2 Suppl):S49-283.

*McNab D, et al. An open label, single-centre, randomized trial of spinal cord stimulation vs. percutaneous myocardial laser revascularization in patients with refractory angina pectoris: the SPiRiT trial. Eur Heart J 2006 May;27(9):1048-53.

Mekhail NA, et al. Retrospective review of 707 cases of spinal cord stimulation: indications and complications. Pain Pract 2011 Mar;11(12):148-53.

Muhammad S, et al. Burst or high-frequency (10 kHz) spinal cord stimulation in failed back surgery syndrome patients with predominant back pain: one year comparative data. Neuromodulation 2017 May 24. [Epub ahead of print].

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 11 of 13

- *National Institute of Clinical Excellence (NICE). Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin. 2008 Oct [www.nice.org] accessed 7/18/17.
- *North RB, et al. Spinal cord stimulation for chronic pain of spinal origin: a valuable long-term solution. Spine 2002 Nov 15; 27(22):2584-91.
- *North RB, et al. Spinal cord stimulation for axial low back pain: a prospective, controlled trial comparing dual and single percutaneous electrodes. Spine 2005 Jun 15;30(12):1412-8.
- *North RB, et al. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a cost effectiveness and cost utility analysis based on a randomized, controlled trial. Neurosurgery 2007 Aug;61(2):361-8.
- Pan X, Bao H, Si Y, et al. Spinal cord stimulation for refractory angina pectoris: a systematic review and meta-analysis. Clin J Pain. Nov 21 2016. [Epub ahead of print].
- *Pedrini L, et al. Spinal cord stimulation for lower limb ischemic pain treatment. Interact Cardiovasc Thorac Surg 2007 Aug;6(4):495-500.
- Peng L, et al. Spinal cord stimulation for cancer-related pain in adults. Cochrane Database Syst Rev 2015 June 29;(6):CD009389.
- Perryman L, et al. Injectable spinal cord stimulation system: A pilot study. Tech Reg Anesth Pain Manag 2012 Feb;16(2):102-5. [http://www.techreganesth.org/article/S1084-208X(13)00011-6/abstract].accessed 7/18/17.
- Perruchoud C, et al. Analgesic efficacy of high-frequency spinal cord stimulation: a randomized double-blind placebo-controlled study. Neuromodulation 2013 July-Aug;16(4):363-369.
- Russo M and Van Buyten JP. 10-kHz high frequency SCS therapy: a clinical summary. Pain Med 2015 May;16(5):934-942.
- Russo M, et al. High frequency spinal cord stimulation at 10 kHz for the treatment of chronic pain: 6-month Australian clinical experience. Pain Physician 2016 May;19(4):267-280.
- Schu S, et al. A prospective, randomized, double-blind, placebo-controlled study to examine the effectiveness of burst spinal cord stimulation patterns for the treatment of failed back surgery syndrome. Neuromodulation 2014 Jul;17(5):443-50.
- Schu S, et al. Spinal cord stimulation of the dorsal root ganglion for groin pain- a retrospective review. Pain Pract 2015 April;15(4):293-299.
- *Sears NC, et al. Long-term outcomes of spinal cord stimulation with paddle leads in the treatment of complex regional pain syndrome and failed back surgery syndrome. Neuromodulation 2011 Jul-Aug;14(4):312-8.
- *Sgueglia GA, et al. Long-term follow-up of patients with cardiac syndrome X treated by spinal cord stimulation. Heart 2007 May;93(5):591-7.
- *Simpson EL, et al. Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin: systematic review and economic evaluation. Health Tech Assess 2009 Mar;13(17):iii, ix-x,1-154.
- Slangen R, et al. Spinal cord stimulation and pain relief in painful diabetic peripheral neuropathy: a prospective two-center randomized controlled trial. Diabetes Care Nov 2014; 37(11):3016-24.
- *Taylor RS, et al. Spinal cord stimulation for chronic back pain and leg pain and failed back surgery syndrome: a systemic review and analysis of prognostic factors. Spine 2005 Jan 1;30(1):152-60.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 12 of 13

*Taylor RS. Spinal cord stimulation in complex regional pain syndrome and refractory neuropathic back and leg pain/failed back surgery syndrome: results of systematic review and meta-analysis. J Pain Symptom Manag 2006 Apr;3(4 Suppl):S13-9.

*Taylor RS, et al. Spinal cord stimulation in the treatment of refractory angina: systematic review and meta-analysis of randomized controlled trials. BMC Cardiovasc Disord 2009 Mar 25;9:13.

Taylor RS, et al. The cost-effectiveness of spinal cord stimulation in the treatment of failed back surgery syndrome. Clin J Pain 2010 Jul-Aug;26(6):463-9.

Taylor RS, et al. Predictors of pain relief following spinal cord stimulation in chronic back and leg pain and failed back surgery syndrome: A systematic review and meta-regression analysis. Pain Pract 2014 Jul;14(6):489-505.

Tiede J, et al. Novel spinal cord stimulation parameters in patients with predominant back pain. Neuromodulation 2013 July-Aug;16(4):370-375.

Torre-Amione G, et al. Spinal cord stimulation is safe and feasible in patients with advanced heart failure: early clinical experience. Eur J Heart Fail Jul 2014;16(7):788-95.

Tsigardas N, et al. Spinal cord stimulation in refractory angina. A systemic review of randomized controlled trials. Acta Cardiol 2015 April;70(2):233-243.

*Ubbink DT, et al. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. Cochrane Database Syst Rev. 2013 Feb 28;2:CD004001.

*Ubbink DT, et al. Spinal cord stimulation for critical leg ischemia: a review of effectiveness and optimal patient selection. J Pain Symptom Manag 2006 Apr;31(4 Suppl):S30-5.

*Vallejo R, et al. Neuromodulation of the cervical spinal cord in the treatment of chronic intractable neck and upper extremity pain: a case series and review of the literature. Pain Physician 2007 Mar;10(2):305-11.

van Beek M, et al. Sustained treatment effect of spinal cord stimulation in painful diabetic peripheral neuropathy: 24-month follow-up of a prospective two-center randomized controlled trial. Diabetes care 2015 Sept;38(9):e132-134.

Van Buyten JP, et al. High-frequency spinal cord stimulation for the treatment of chronic back pain patients: results of a prospective multicenter European clinical study. Neuromodulation 2013 Jan-Feb;16(1):59-65.

Van Buyten JP, et al. stimulation of the dorsal root ganglia for the management of complex regional pain syndrome: a prospective case series. Pain Pract 2015 March;15(3):208-216.

Weiner RL, et al. Treatment of failed low back pain with a novel percutaneous DRG wireless stimulator: pilot and feasibility study. Pain med 2016 Oct;17(10):1911-1916.

Williams KA, et al. A multi-center analysis evaluating factors associated with spinal cord stimulation outcome in chronic pain patients. Pain Med 2011 Aug;12(8):1142-53.

Wolter T, et al. The impact of psychological factors on outcomes for spinal cord stimulation: an analysis with long-term follow-up. Pain Physician 2013 May-Jun;16(3):265-75.

Wolter T, et al. Cervical spinal cord stimulation: an analysis of 23 patients with long-term follow-up. Pain Physician 2012 May-Jun;15(3):203-12.

Zipes DP, et al. Spinal cord stimulation therapy for patients with refractory angina who are not candidates for revascularization. Neuromodulation 2012 Nov-Dec;15(6):550-8.

Zipes DP, et al. Determining the feasibility of spinal cord neuromodulation for the treatment of chronic systolic heart failure: the DEFEAT-HF study. JACC Heart Fail 2016 Feb;4(2):129-136

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 13 of 13

*Key Article

KEY WORDS

Burst stimulation, Dorsal column, Dorsal root ganglion, High-frequency neurostimulation, Neuromodulation, Neurostimulation, Wireless neurostimulation.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for electrical nerve stimulators that includes dorsal column stimulators. Please refer to the following NCD website for Medicare Members:

<http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=240&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&CptHcpcsCode=36514&bc=gAAAABAAAA&.>