

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	LUMBAR DECOMPRESSION
Policy Number	7.01.97
Category	Technology Assessment
Effective Date	06/21/18
Revised Date	12/20/18
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based on our criteria and assessment of the peer-reviewed literature, an initial primary lumbar decompression has been medically proven to be effective and is considered **medically appropriate** for spinal stenosis/spondylolisthesis when ALL the following are met:
 - A. Subjective symptoms including at least **TWO** of the following:
 1. Significant level of pain on a daily basis defined on a Visual Analog Scale (VAS/Numeric Rating Scale (NRS) as greater than or equal to 7;
 2. Persistent radiating pain into the buttock(s) and/or lower extremity(ies) on a daily basis that has a documented negative impact on activities of daily living despite optimal conservative treatment as described below:
 3. Pain, cramping, weakness, or tingling in the lower back, buttock(s), and leg(s) brought about by walking or positions that cause thecal sac or nerve root compression (e.g., standing, extension)
 - B. Performed for **EITHER** of the following:
 1. Neurogenic claudication secondary to central/lateral recess/foraminal stenosis when ALL of the following criteria are met:
 - a. Subjective symptoms including **EITHER** of the following:
 - i. Symptoms worsen with standing and/or walking; or
 - ii. Symptoms are alleviated with sitting and/or forward flexion.
 - b. Objective physical findings consistent with recent (within 6 months) MRI/CT; and
 - c. Less than clinically meaningful improvement from epidural steroid injections/selective nerve root block(s) unless contraindicated.
 2. Spondylolisthesis with neurogenic claudication symptoms or radicular pain from lateral recess, or foraminal stenosis associated with Listhesis demonstrated on plain x-rays.
 - C. Less than clinically meaningful improvement with **EITHER** of the following unless contraindicated:
 1. Prescription strength analgesics, steroids, and/or NSAIDS for 6 weeks; or
 2. Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician for 6 weeks.
 - D. Recent (within 6 months) MRI/CT identifies nerve root impingement and/or thecal sac impingement caused by stenosis/listhesis that correlates with patient symptoms or physical findings.
 - E. No previous surgeries at the level(s) involved.
 - F. All other sources of pain have been excluded.

Medical Policy: LUMBAR DECOMPRESSION

Policy Number: 7.01.97

Page: 2 of 5

- G. Absence of unmanaged significant behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, drug and alcohol abuse).
- II. Based on our criteria and assessment of the peer-reviewed literature, a repeat lumbar decompression has been medically proven to be effective and is considered **medically appropriate** when ALL of the criteria noted in Policy Statement I above (with the exception of E, no previous surgeries at level(s) involved) plus ALL of the following are met:
 - A. Recent (within 3 months) post-operative MRI with gadolinium/CT myelogram confirms radiographic evidence of neural structure compression;
 - B. Greater than 12 months since last decompression surgery; and
 - C. Initial relief of symptoms following previous decompression procedure at same level(s).
- III. Based upon our criteria and assessment of peer-reviewed literature, the following procedures have not been medically proven effective and are therefore considered **investigational** or unproven:
 - A. Percutaneous lumbar discectomy;
 - B. Percutaneous laser discectomy;
 - C. Laser-assisted disc decompression; and
 - D. Percutaneous nucleotomy.
- IV. Based upon our criteria and assessment of peer-reviewed literature, Interspinous/interlaminar process spacer devices (ISS) and interspinous/interlaminar stabilization/distraction devices, and interspinous process decompression (IPD) systems/devices (e.g., Coflex Interlaminar Technology Implant, Superior ISS Interspinous Spacer System, X-STOP PEEK Interspinous Process Decompression System) have not been medically proven effective and are therefore considered **investigational** and/or unproven for ALL the following indications including, but not limited to:
 - A. Lumbar interspinous/interlaminar distraction without fusion for indirect spinal decompression;
 - B. Lumbar interspinous fixation with fusion with or without decompression for stabilization;
 - C. Lumbar spinal stabilization with an interspinous process device/interlaminar device without fusion in conjunction with decompression laminectomy.

POLICY GUIDELINES

- I. Acceptable imaging modalities are CT scan, MRI and myelogram. Imaging must be performed and read by an independent radiologist. If discrepancies should arise in the interpretation of the imaging, interpretations by the radiologist will supersede. Discography results will not be used as a determining factor of medical necessity for any requested procedures. Use is not endorsed.
- II. Clinically meaningful improvement is defined as global assessment showing at least 50% improvement

DESCRIPTION

Narrowing/stenosis or spondylolisthesis that creates a narrowing of the spinal canal can cause chronic pain, numbness, and muscle weakness in an individual's arms or legs. Spinal decompression can be performed anywhere along the spine from the neck (cervical) to the lower back (lumbar). The procedure is performed through a surgical incision in the back (posterior). The lamina is the bone that forms the backside of the spinal canal and makes a roof over the spinal cord. Removing the lamina and other soft tissues gives more room for the nerves, relieves pressure and allows for removal of bone spurs. Depending on the extent of stenosis, one vertebra (single-level) or more (multi-level) may be involved. There are several types of decompression surgery:

- I. Laminectomy is the removal of the entire bony lamina, a portion of the enlarged facet joints, and the thickened ligaments overlying the spinal cord and nerves.
- II. Laminotomy is the removal of a small portion of the lamina and ligaments, usually on one side. Using this method the natural support of the lamina is left in place, decreasing the chance of postoperative spinal instability. Sometimes an endoscope may be used, allowing for a smaller, less invasive incision.

Medical Policy: LUMBAR DECOMPRESSION

Policy Number: 7.01.97

Page: 3 of 5

III. Foraminotomy is the removal of bone around the neural foramen - the space between vertebrae where the nerve root exits the spinal canal. This method is used when disc degeneration has caused the height of the foramen to collapse, resulting in a pinched nerve. It can be performed with a laminectomy or laminotomy.

RATIONALE

The Spine Patient Outcomes Research Trial (SPORT) was funded by the National Institutes of Health (NIH) to study the outcomes from surgical and nonsurgical management of three conditions: intervertebral disc herniation, degenerative spondylolisthesis, and lumbar spinal stenosis. Both surgical and nonsurgical care of intervertebral disc herniation resulted in significant improvement in symptoms of low back and leg pain. However, the treatment effect of surgery for intervertebral disc herniation was less than that seen in individuals with degenerative spondylolisthesis and lumbar spinal stenosis. The preliminary 4-year outcomes data demonstrated more significant degrees of improvement in pain levels and function with surgical versus nonsurgical treatment in the chronic conditions of lumbar spinal stenosis and lumbar spinal stenosis with spondylolisthesis (Asghar, 2012; Weinstein, 2006a; Weinstein, 2006b; Weinstein, 2007; Weinstein, 2009).

According to the American Pain Society (APS), decompressive laminectomy may be an acceptable option for individuals experiencing disabling and persistent leg pain due to spinal stenosis, either with or without degenerative spondylolisthesis. The APS reports that decompressive laminectomy is associated with moderate benefits compared to nonsurgical therapy through 1 to 2 years, though the effects of the procedure appear to diminish with long-term follow-up. Although individuals on average do not worsen without surgery, improvements are less than those observed in individuals with radiculopathy due to herniated lumbar disc. Their guidelines indicate there is insufficient evidence to determine if laminectomy with fusion is more effective than laminectomy without fusion. The authors recommended that shared decision-making regarding surgery include a specific discussion about moderate/average benefits, which appear to decrease over time in affected individuals who undergo surgery (Chou, 2009).

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
22867 (E/I)	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level
22868 (E/I)	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (List separately in addition to code for primary procedure)
22869 (E/I)	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level
22870 (E/I)	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (List separately in addition to code for primary procedure)
63005	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (eg, spinal stenosis), 1 or 2 vertebral segments; lumbar, except for spondylolisthesis

Medical Policy: LUMBAR DECOMPRESSION**Policy Number: 7.01.97****Page: 4 of 5**

Code	Description
63011	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (eg, spinal stenosis), 1 or 2 vertebral segments; sacral
63012	Laminectomy with removal of abnormal facets and/or pars inter-articularis with decompression of cauda equina and nerve roots for spondylolisthesis, lumbar (Gill type procedure)
63017	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (eg,
63047	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [eg, spinal or lateral recess stenosis]), single vertebral segment; lumbar
63048	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [eg, spinal or lateral recess stenosis]), single vertebral segment; each additional segment, cervical, thoracic, or lumbar

Copyright © 2018 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No codes	

ICD10 Codes

Code	Description
M43.10-M43.17	Spondylolisthesis, thoracolumbar, lumbar or lumbosacral region (code range)
M48.05-M48.07	Spinal stenosis, thoracolumbar, lumbar or lumbosacral region (code range)

REFERENCES

Andreisek G, et al. Uncertainties in the diagnosis of lumbar spinal stenosis. Radiology 2011;261(3):681-4.

Backstrom KM, et al. Lumbar spinal stenosis-diagnosis and management of the aging spine. Manual Therapy 2011;16(4):308-17.

Bae HW, et al. Three-year follow-up of the prospective, randomized, controlled trial of corflex interlaminar stabilization vs. instrumented fusion in patients with lumbar stenosis. Neurosurgery 2016;79(2):169-81.

Bae HW, et al. Therapeutic sustainability and durability of cofex interlaminar stabilization after decompression for lumbar spinal stenosis: a four year assessment. International Journal of Spine Surgery. 2015;9.

Brox JI, et al. Four-year follow-up of surgical versus non-surgical therapy for chronic low back pain. Annals of the Rheumatic Diseases 2010;69(9):1643-8.

Choi D, et al. Review of metastatic spine tumour classification and indications for surgery: the consensus statement of the Global Spine Tumour Study Group. European Spine Journal 2010;19(2):215-22.

Chou R, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. Annals of Internal Medicine 2007;147(7):478-91.

Chou R, et al. Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/ American College of Physicians clinical practice guideline. Annals of Internal Medicine 2007;147(7):505-14.

Medical Policy: LUMBAR DECOMPRESSION

Policy Number: 7.01.97

Page: 5 of 5

Chou R, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. Spine 2009;34(10):1066-77.

Conn A, et al. Systematic review of caudal epidural injections in the management of chronic low back pain. Pain Physician 2009;12(1):109-35.

Darouiche RO. Spinal epidural abscess. New England Journal of Medicine 2006;355(19):2012-20.

Deyo RA, et al. Trends, major medical complications, and charges associated with surgery for lumbar spinal stenosis in older adults. Journal of the American Medical Association 2010;303(13):1259-65.

Eliyas JK, et al. Surgery for degenerative lumbar spine disease. Disease-a-Month 2011;57(10):592-606.

Fu KM, et al. Morbidity and mortality in the surgical treatment of 10,329 adults with degenerative lumbar stenosis. Journal of Neurosurgery: Spine 2010;12(5):443-6.

Hahne AJ, et al. Conservative management of lumbar disc herniation with associated radiculopathy: a systematic review. Spine 2010;35(11):E488-504.

Kalichman L, Hunter DJ. Diagnosis and conservative management of degenerative lumbar spondylolisthesis. European Spine Journal 2008;17(3):327-35.

Kanayama M, et al. Effective prevention of surgical site infection using a Centers for Disease Control and Prevention guideline-based antimicrobial prophylaxis in lumbar spine surgery. Journal of Neurosurgery: Spine 2007;6(4):327-9.

Klein G, et al. Nonoperative treatment of spondylolysis and grade I spondylolisthesis in children and young adults: a meta-analysis of observational studies. Journal of Pediatric Orthopedics 2009;29(2):146-56.

Kreiner DS, et al. Diagnosis and treatment of degenerative lumbar spinal stenosis. Evidence-based clinical guidelines for multidisciplinary spine care. North American Spine Society. 2011. [<http://www.spine.org>]. accessed 2/19/18.

Sansur CA, et al. Morbidity and mortality in the surgical treatment of 10,242 adults with spondylolisthesis. Journal of Neurosurgery: Spine 2010;13(5):589-93.

Selective dorsal rhizotomy for spasticity in cerebral palsy. NICE interventional procedure guidance IPG373 National Institute for Health and Clinical Excellence. 2010 Dec. [<http://guidance.nice.org.uk>]. accessed 2/19/18.

Thome C, et al. Outcome after less-invasive decompression of lumbar spinal stenosis: a randomized comparison of unilateral laminotomy, bilateral laminotomy, and laminectomy. Journal of Neurosurgery: Spine 2005;3(2):129-41.

Tran de QH, Duong S, Finlayson RJ. Lumbar spinal stenosis: a brief review of the nonsurgical management. Canadian Journal of Anaesthesia 2010;57(7):694-703.

*Key Article

KEY WORDS

Lumbar foraminotomy, Lumbar decompression, Lumbar laminectomy, Spinal stenosis, Spondylolisthesis

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based upon review, lumbar decompression is not addressed in a National or Local Medicare coverage determination or policy