

# MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	SURGICAL MANAGEMENT OF OBESITY
Policy Number	7.01.29
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	03/21/02, 02/20/03, 10/15/03, 11/18/04, 08/18/05, 04/20/06, 11/16/06, 11/15/07, 12/18/08, 01/21/10, 12/16/10, 10/20/11, 12/20/12, 11/21/13, 12/18/14, 11/19/15, 10/20/16, 4/20/17, 03/15/18, 08/16/18, 05/16/19
Product Disclaimer	<ul style="list-style-type: none"> <li>• If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</li> <li>• If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</li> <li>• If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</li> </ul>

## POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, the surgical treatment of morbid obesity by *open or laparoscopic Roux-en-Y gastric bypass, duodenal switch procedure (biliopancreatic diversion), and sleeve gastrectomy* have been medically proven to improve health outcomes and are therefore **medically appropriate** for selected patients. Patients must meet all of the criteria:
- A. Patients must be morbidly obese, which is defined as:
1. *For adults, ages 18 and older*: either having a BMI greater than or equal to 40 kg/m<sup>2</sup> or having a BMI greater than or equal to 35 kg/m<sup>2</sup> and existing comorbid condition(s) (e.g., hypertensive cardiovascular disease, coronary heart disease, pulmonary hypoventilation, hypertension, hypercholesterolemia, dyslipidemias, diabetes, sleep apnea, degenerative arthritis of weight-bearing joints, or other weight related arthropathies, and metabolic syndrome). Documentation of the comorbid existing medical condition(s) must be submitted by the primary care physician; or
  2. *For adolescents, ages 16-18 years old*: either having a BMI greater than or equal to 40 kg/m<sup>2</sup> (greater than 140% of the 95th percentile) or having a BMI greater than or equal to 35 kg/m<sup>2</sup> (greater than 120% of the 95th percentile) and existing comorbid condition(s) (e.g., hypertensive cardiovascular disease, hypertension, hypercholesterolemia, dyslipidemias, diabetes, sleep apnea, weight related arthropathies, and metabolic syndrome). Documentation of the comorbid existing medical condition(s) must be submitted by the primary care physician.
- B. A letter of support from the physician currently providing primary care to the member and who is familiar with his/her attempts at weight reduction, medical history and current health status (including obesity issues) is also necessary for the review process.
- C. History of rigorous attempts at weight reduction.
1. There must be written evidence of a weight loss history, either by the bariatric surgeon, primary care physician or nutritionist. This documentation should include the name of the weight loss program, length of participation in the program and any weight loss achieved. At least one program must have been a supervised weight loss program;
  2. Weight loss attempts need not be continuous, but a minimum total of six (6) months is required.
    - a. If the patient has had no previous attempts at medical weight loss, participation for a minimum of six (6) months in a preoperative bariatric surgery weight loss program is required.

## Medical Policy: SURGICAL MANAGEMENT OF OBESITY

Policy Number: 7.01.29

Page: 2 of 17

- b. If past prior attempts of weight loss are remote (greater than 5 years), then the patient is required to participate, for a minimum of six (6) months in a preoperative bariatric surgery weight loss program.
  - D. There must be no significant liver, kidney, or gastrointestinal disease present. The presence of non-alcoholic steatohepatitis or “fatty liver”, which is associated with morbidly obese patients, would not be considered significant liver disease in this instance.
  - E. Treatable metabolic causes for obesity (e.g., adrenal or thyroid disorders) have been addressed.
  - F. Patients with a history of alcohol or substance abuse will not be considered unless there is a record of at least six months of abstinence. If there has been six months of abstinence, this condition must be addressed in a psychiatric consultation.
  - G. Patients must be screened by their physician for major psychopathology. All patients who have current symptoms which concern the physician, or who have had a psychiatric hospitalization must have a psychiatric *evaluation*. The psychiatric evaluation *should* be performed by a psychiatrist familiar with the implications of weight reduction surgery. If psychiatrists with this expertise are not available, an evaluation by a clinical psychologist familiar with the implications of weight reduction surgery is also acceptable. A psychiatrist or clinical psychologist who is providing ongoing care for the patient may also provide this evaluation. If the patient already has an established psychiatric provider, that provider (e.g., psychiatrist or clinical psychologist) must provide a second letter of support for the proposed surgery. Psychological *testing* as screening tool or as part of the psychological evaluation prior to bariatric surgery is considered **not medically necessary**.
- II. Based upon our criteria and assessment of peer-reviewed literature, the surgical management of morbid obesity by *laparoscopic adjustable gastric banding* (e.g., LAP BAND) is considered **medically appropriate** in the following circumstances:
- A. The patient must meet all the requirements listed above in Policy Statement I, A-G; and
  - B. The dietary history does not include a large consumption of high caloric liquids (e.g., milk shakes) or sweets; and
  - C. The patient has no significant history of esophageal or gastric disease (*please note contraindications to adjustable gastric banding listed in the rationale section*); and
  - D. The patient must participate in a pre-operative bariatric program that requires a 5% weight loss to demonstrate commitment to behavioral and dietary changes. The 5% weight loss will be measured from the date of the patient’s initial visit to the bariatric surgeon to the date of the request for pre-authorization of the adjustable gastric banding procedure.
- III. Based upon our criteria and assessment of peer reviewed literature, although the supportive literature is limited, surgical procedures for the management of obesity for *patients between 16-18* will be considered **medically necessary** if the patient meets all the criteria in Policy Statement I including history of rigorous attempts at weight reduction as noted in Policy statement I. C. of at least one year for this age group.
- IV. Based upon our criteria and assessment of peer reviewed literature, there is insufficient information to support use of surgical procedures for the management of obesity for *patients under age 16*. Therefore, surgery for obesity is considered **investigational** for this age group.
- V. Based upon our criteria and assessment of the peer-reviewed literature, the following procedures for the primary surgical treatment of morbid obesity have not been medically proven to improve health outcomes and therefore are considered **investigational**:
- A. Aspiration therapy (e.g., AspireAssist device);
  - B. Laparoscopic gastric plication;
  - C. Mini-gastric bypass (also called loop gastric by-pass);
  - D. Single anastomosis duodenoileal bypass with sleeve gastrectomy (SADI-S);
  - E. Stomach intestine pylorus sparing surgery (SIPS); and
  - F. Transoral gastroplasty or endoscopic/endoluminal procedures (e.g., Restorative Obesity Surgery, Endoluminal or ROSE, TOGA System, StomaphyX, EndoCinch, Overstitch device, EndoBarrier, intragastric balloon insertion).
- VI. *Surgical revisions* are considered **medically appropriate** for complications, such as malabsorption/malnutrition, obstruction, staple disruption, or stricture following the primary procedure. (*Please see Guidelines section, #IV*).

## Medical Policy: SURGICAL MANAGEMENT OF OBESITY

Policy Number: 7.01.29

Page: 3 of 17

- VII. A *revision or removal of a laparoscopic adjustable gastric band* is considered **medically necessary** for complications or for a technical failure. Examples of laparoscopic adjustable gastric banding complications reported in the literature that may warrant revision, removal or conversion to another procedure include, but are not limited to, band slippage, band erosion, infection, esophageal dilation, dysphagia, and heartburn/reflux. Technical failures of LAGB include (but are not limited to) a displaced band, port dislocation, too tight a band creating food passage problems, band intolerance (e.g., pain, vomiting), and port and/or catheter leakage.
- VIII. A *revision or conversion* to another medically appropriate procedure because of unsatisfactory weight loss due to technical failure of the primary bariatric procedure such as pouch dilation or an initial pouch size that is too large (an ideal initial pouch size is approximately 20 cc) is considered **medically appropriate** if there is documentation regarding all of the following submitted with the request:
- A. Primary procedure was initially successful in inducing weight loss; and
  - B. Patient has remained compliant for at least six months to the prescribed nutrition and exercise program. (*See Guidelines section #III*).
- IX. Placement of a *second* adjustable gastric band is considered **investigational** as there is no published literature to support the efficacy of a second adjustable gastric band after failure to produce weight loss of the first procedure.
- X. Based upon our criteria and assessment of peer-reviewed literature, use of an endoscopic/endoluminal procedure for *revisional* surgery (e.g., transoral outlet reduction) has not been medically proven effective and is considered **investigational**.
- XI. *Repeat surgery* for morbid obesity is considered **not medically necessary** for those patients who have either failed to lose weight or who have regained weight due to non-adherence with the prescribed nutrition and exercise program following their surgery.
- XII. Based upon our criteria and assessment of peer-reviewed literature, use of bariatric surgery as a treatment for either *non-obese patients with* Type 2 diabetes mellitus or for Type 2 diabetics with BMI between 30.0-34.9 kg/m<sup>2</sup> has not been medically proven to be effective and is considered **investigational**.
- XIII. Based upon our criteria and assessment of peer-reviewed literature, performing a *routine liver biopsy* at the time of the bariatric surgery is considered **not medically necessary** in the absence of objective signs or symptoms of liver disease (e.g., elevated liver enzymes, enlarged liver).
- XIV. Based upon our criteria and assessment of peer-reviewed literature, prophylactic removal of a normal and asymptomatic gallbladder at the time of the bariatric surgery is considered **not medically necessary** unless cholelithiasis is present or for patients who will undergo biliopancreatic diversion based on a higher incidence of biliary complications.
- XV. The adjustment of a previously placed laparoscopic gastric band, beyond the global, 90-day limit, is considered medically necessary to control the rate of weight loss and/or treat symptoms secondary to gastric restriction following a **medically necessary** adjustable gastric banding procedure.

*Refer to Corporate Medical Policy #11.01.03 regarding Experimental and Investigational Services.*

*Refer to Corporate Medical Policy # 11.01.11 regarding Cosmetic and Reconstructive procedures.*

*Refer to Corporate Medical Policy # 7.01.05 regarding Vagus Nerve Stimulation and Vagus Nerve Blocking Therapy.*

*Refer to Corporate Medical Policy # 7.01.64 regarding Gastric Electrical Stimulation.*

### **POLICY GUIDELINES**

- I. Patients considering surgery must participate in an integrated pre- and post-surgery program consisting of dietary therapy, physical activity, and behavioral and social support programs. Post-surgically, patients must be involved in a formal program for at least one year.
- II. Gastric bypass surgery affects the absorption of medication and may lead to irregular blood levels of medication. Where drug level maintenance is critical, bypass surgery may be contraindicated. Examples include patients with

## Medical Policy: SURGICAL MANAGEMENT OF OBESITY

Policy Number: 7.01.29

Page: 4 of 17

seizure disorders requiring anti-seizure medications and patients with mental illness who require maintenance medication.

- III. Some post bariatric surgery patients regain lost weight or never lose sufficient weight. Other patients may develop unacceptable postoperative symptoms. These failures may warrant reversal surgery or revision surgery (e.g., conversion to Roux-en-Y). Failures due to patient noncompliance reflect poor patient selection and do not warrant revision procedures. A clue to this is gastric pouch dilation in a patient not adhering to the recommended eating protocols. These patients are likely to fail again. Patients must demonstrate compliance for at least six months before a revision will be considered for failure to lose adequate weight and/or weight regain. Revision due to medical complications does not require six months of demonstrated compliance.
- IV. Coverage is limited to physicians who have been properly trained in performing a bariatric procedure at facilities with the diagnostic and support services necessary for the care of morbidly obese patients.
- V. An expected outcome of successful bariatric surgery is redundant/excessive skin. Surgery to remove this skin is generally not considered medically necessary and therefore not covered. *Refer to Corporate Medical Policy # 7.01.53 regarding Abdominoplasty and panniculectomy.*
- VI. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

### **DESCRIPTION**

Obesity is a complex multifactorial, chronic condition that substantially raises an individual's risk of morbidity from hypertension, dyslipidemia, type 2 diabetes, coronary artery disease, stroke, gall bladder disease, osteoarthritis, sleep apnea, respiratory problems and a variety of cancers.

The 1998 National Heart Lung Blood Institute (NHLBI) expert panel defined Class I obesity as a body mass index (BMI) of 30-34.9 kg/m<sup>2</sup>; Class II obesity as a BMI of 35-39.9 kg/m<sup>2</sup>; and extreme obesity or Class III as a BMI greater than or equal to 40 kg/m<sup>2</sup>.

Bariatric surgery can be divided into two categories: gastric restrictive procedures and malabsorptive procedures. Gastric restrictive procedures mechanically prevent the patient from overeating and the latter procedures interfere with the absorption of ingested nutrients. Examples of gastric restrictive procedures include vertical and horizontal banded gastroplasty and adjustable gastric banding. Malabsorptive procedures include biliopancreatic bypass, long limb gastric bypass and biliopancreatic bypass with duodenal switch. The Roux-en-Y gastric bypass is actually a combination of a gastric restrictive and malabsorptive procedure. A bariatric procedure is usually considered a success when at least 50% of excess body weight is lost or when the patient returns to within 30% of his/her ideal body weight.

The intragastric balloon has been proposed as a temporary non-surgical obesity treatment for short-term weight loss in patients who have had unsatisfactory results with their diet and exercise programs. The intragastric balloon has also been proposed for weight loss in the super-obese prior to a permanent, invasive surgical procedure. The saline-filled intragastric balloon, placed endoscopically, is intended to reduce gastric capacity, creating satiety and reducing food intake.

The TOGA and StomaphyX Systems consist of a set of transoral endoscopically guided staples that are used to create a stapled restrictive pouch along the lesser curvature of the stomach. The EndoCinch was initially devised for the endoscopic treatment of gastroesophageal reflux disease (GERD). With EndoCinch, sutures are deployed in a continuous and cross-linked fashion from the proximal fundus to the distal body. Once the suture is fixed, distention of the stomach is significantly limited, thus providing a method of restricting food intake. The Overstitch™ device allows for a full thickness endoscopic suturing compared to superficial thickness suturing provided by other devices. These endoscopic procedures may offer lower morbidity than other current bariatric procedures and are adjustable and reversible. They are being investigated as the primary bariatric surgery and as a revisional procedure to treat weight gain (e.g., large gastric pouch, large gastric stoma/dilated gastrojejunal anastomosis).

Laparoscopic gastric plication is a relatively new restrictive technique that involves sewing one or more folds in the stomach. Gastric plication reduces stomach volume by approximately 70%, and is potentially reversible.

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 5 of 17**

Aspiration therapy as a treatment for obesity involves a surgically placed tube to drain a portion of the stomach contents after every meal. The tube is inserted in the stomach with an endoscope via a small incision in the abdomen. A disk-shaped port valve that lies outside the body, flush against the skin of the abdomen, is connected to the tube and remains in place. Approximately 20 to 30 minutes after meal consumption, the patient attaches the device's external connector and tubing to the port valve, opens the valve and drains the contents. Once opened, it takes approximately five to 10 minutes to drain food matter through the tube and into the toilet. The device removes approximately 30 percent of the calories consumed.

Single anastomosis duodeno-ileal bypass with sleeve gastrectomy (SADI-S) has a restrictive component when reducing the greater curvature of the stomach, but especially a malabsorptive component, as the common channel is also reduced. The objective of this surgical technique is to lessen the intestinal loop where nutrients are absorbed. The procedure is based on the biliopancreatic diversion in which a sleeve gastrectomy is followed by an end-to-side duodeno-ileal diversion. The preservation of the pylorus makes possible the reconstruction in one loop.

Stomach intestine pylorus sparing surgery (SIPS) is a modified version of the duodenal switch procedure. SIPS involves the creation of a 300 cm common channel with a single-anastomosis duodenal enterostomy. It involves the formation of a sleeve gastrectomy that is slightly larger than our usual sleeve, with an attachment placed beneath the pyloric valve, which controls emptying of the stomach into the mid gut, located three meters from the terminal ileum. A proposed benefit of SIPS is that it does not cause abrupt rise and fall of blood glucose, thus preserving the pyloric valve. Also, by not bypassing as much intestine, it may reduce the complications of short bowel syndrome.

Post bariatric surgery patients who regain lost weight, do not lose sufficient weight or develop unacceptable postoperative symptoms due to structural complications may warrant reversal or revision surgery. Reversal or revision of bariatric procedures is usually not warranted in patient failure due to noncompliance (e.g. gastric pouch dilation from 20cc to greater than 100cc in a patient not adhering to the recommended eating protocols). Conversely, revisional surgery for complications related to malabsorption resulting in hypoglycemia, malnutrition or weight loss of 20% below ideal body weight have been noted, particularly after malabsorptive procedures. Examples of laparoscopic adjustable gastric banding complications reported in the literature that may warrant revision, removal or conversion to another procedure include, but are not limited to, band slippage, band erosion, infection, esophageal dilation, dysphagia, and heartburn/reflux. Technical failures of LAGB include (but are not limited to) a displaced band, port dislocation, too tight a band creating food passage problems, band intolerance (e.g., pain, vomiting), and port and/or catheter leakage.

### **RATIONALE**

The hallmark piece of literature published supporting the safety and effectiveness of bariatric surgery is the National Institutes of Health (NIH) Consensus Statement. Among their findings, the panel recommended (1) gastric restrictive or bypass procedures could be considered for well informed and motivated individuals with acceptable operative risks (2) patients who are candidates for surgical procedures should be selected carefully after evaluation by a multidisciplinary team with medical, surgical, psychiatric and nutritional expertise, (3) the operation be performed by a surgeon substantially experienced with appropriate procedures and working in a clinical setting with adequate support for all aspects of management and assessment and (4) lifelong medical surveillance after surgical therapy is necessary.

The NIH Conference on the Surgical Management of Morbid Obesity (1998) states that obesity surgery should be reserved only for patients who have first attempted medical therapy. The NIH Consensus Conference states that the initial goal of medical therapy is a 10% weight reduction with a reasonable duration of six months. "The rationale for this initial goal is that even moderate weight loss can significantly decrease the severity of obesity-associated risk factors." The patient's ability to lose weight prior to bariatric surgery is an indication of the likelihood of compliance with the severe dietary restriction and behavioral changes required for the patient following surgery. Weight loss prior to surgery can also make surgical intervention easier to perform.

The Swedish Obese Subjects (SOS) intervention trial has reported on several hundred patients with up to 8 years of follow up. This trial demonstrated that surgery results in large amounts of weight loss compared to usual care (16% decrease in total body weight at six years versus an increase of 0.8% for usual care). Results of the SOS trial show substantial weight loss in the surgery group of a peak of 44 kg at one year and a gradual weight increase thereafter (mean loss of 30 kg at two years and 20 kg at eight years). The SOS trial also shows that co-morbid conditions and quality of life are improved

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 6 of 17**

after surgery, with the most compelling evidence in the co-morbid conditions that exist for diabetics (a 18.5% decrease in diabetes vs. 3.6%). A decrease in the number of patients presenting with hypertension decreased in the short-term, but was not sustained two years following surgery.

In regards to specific bariatric procedures, there is sufficient data published in the medical literature to conclude that surgical management of obesity using open or laparoscopic Roux-en-Y gastric bypass procedures, sleeve gastrectomy or the duodenal switch procedure improves health outcomes for patients with morbid obesity. Improved health outcomes have been achieved outside the investigational setting.

The FDA has given premarket approval for the LAP-BAND® and Realize™ adjustable gastric banding devices. Adjustable gastric banding has been an evolving procedure with issues of migration and erosion being addressed by varying techniques and surgical modifications. Adjustable gastric banding is associated with fewer early complications, but more late complications and re-operations when compared to RGBP. Studies report reoperation rates due to adverse events at 5.6%-24%. Weight loss is also significantly less than Roux-en-Y gastric bypass in most reported studies of AGB. Evidence for 3-year outcomes related to weight loss after AGB was consistent with a 25% or greater excess weight loss reported. The reduction in BMI ranged from 7.9-15, but the typical patient in the studies remained obese (BMI greater than 30) in the majority of the studies. As studies demonstrate the % EWL is significantly less with adjustable gastric banding, the procedure works best for appropriately selected patients. Contraindications to adjustable gastric banding per the product insert of the LAP-BAND® are as follows:

1. Patients with inflammatory diseases of the gastrointestinal tract, including severe intractable esophagitis, gastric ulceration, duodenal ulceration, or specific inflammation such as Crohn's disease.
2. Patients with severe cardiopulmonary diseases or other serious organic disease which may make them poor surgical candidates.
3. Patients with potential upper gastro-intestinal bleeding such as esophageal or gastric varices or congenital acquired intestinal telangiectases.
4. Patients with portal hypertension.
5. Patients with congenital or acquired anomalies of the GI tract such as atresias or stenoses.
6. Patients who have/experience an intra-operative gastric injury during the implantation procedure, such as a gastric perforation at or near the location of the intended band placement.
7. Patients with cirrhosis.
8. Patients with chronic pancreatitis.
9. Patients who are addicted to alcohol and/or drugs.
10. Non-adult patients (patients under 18 years of age).
11. Patients who have an infection anywhere in their body or where the possibility of contamination prior to or during the surgery exists.
12. Patients on chronic, long-term steroid treatment.
13. Patients who are unable or unwilling to comply with dietary restrictions, which are required by this procedure.
14. Patients who are known to have, or suspected to have, an allergic reaction to materials contained in the system or who have exhibited pain intolerance to implanted devices.
15. Patients or family members with a known diagnosis or pre-existing symptoms of autoimmune connective tissue disease such as systemic lupus erythematosus or scleroderma.
16. Pregnancy: Placement of the LAP-BAND System is contraindicated for patients who currently are or may be pregnant. Patients who become pregnant after band placement may require deflation of their bands."

In February 2011, the FDA granted approval for LAGB in patients with a BMI of 30-35 kg/m<sup>2</sup> in the presence of at least one weight-related comorbidity. The FDA labeling indicates that this procedure should be reserved for patients who are at the highest risk for weight-related complications and who have been unsuccessful in achieving medical weight loss. The evidence is insufficient to determine if LAGB improves the overall health outcomes for patients with BMIs that are lower than the current thresholds for bariatric surgery. The patients in these studies consist of a heterogenous patient population and the number and severity of comorbidities vary considerably. While the short-term evidence in current studies demonstrate weight loss in this patient population, and favorable changes in measures of diabetes, the impact of LAGB on other weight-related comorbidities is less certain.

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 7 of 17**

JB Dixon, et al. (2008) performed a randomized controlled trial designed to determine if surgically induced weight loss results in better glycemic control and less need for diabetes medication than conventional approaches to weight loss and diabetes control in patients with BMI of greater than 30 and less than 40. (Results were not reported separately for patients with BMI less than or greater than 35.) Sixty patients were enrolled and 30 were randomized to LAGB and 30 to conventional diabetes care. Fifty-five completed the 2-year follow-up. Remission of diabetes was achieved by 22 (73%) in the LAGB group and 4 (13%) in the control group. The surgical group lost 62.5% of excess weight (using BMI of 25 as ideal weight) versus a loss of 4.3% of excess weight in the conventional group. Mean hemoglobin A1c was less than 6.2% at baseline in two surgically and four conventionally treated patients versus 24 and six patients, respectively, at two years. At baseline, two surgically treated and four conventionally treated patients were using no pharmacotherapy versus 26 and eight, respectively, at two years. One surgical patient developed a wound infection, two developed gastric pouch enlargement and had laparoscopic revision to remove and replace the band. The remaining evidence at the present time consists of small case series and case reports with short follow-up from non-U.S. centers employing procedures considered investigational in this policy (e.g., stand-alone sleeve gastrectomy, mini gastric bypass). Overall, the data are insufficient to allow conclusions regarding the efficacy of bariatric surgical procedures in the treatment or cure of type 2 diabetes.

A September 2004 report addressing bariatric surgery in morbidly obese adolescents found, of all the patients who receive bariatric surgery, about 1% are adolescents. None of the studies investigating bariatric surgery in this age group reported long-term outcomes. Evidence is insufficient to determine if bariatric surgery allows for a clinically significant weight loss amount nor does the evidence allow any conclusions regarding improvements in co-morbidities, or the rates of harm in performing this type of procedure in this patient population. As yet, we do not know how bariatric surgery affects the growth and sexual development of adolescents with morbid obesity.

In 2018, the American Society for Metabolic and Bariatric Surgery (ASMBS) Pediatric Committee updated their evidence-based guidelines published in 2012. The committee identified a significant increase in data supporting the use of metabolic and bariatric surgery (MBS) in adolescents since 2012. ASMBS recommends adolescents with BMI  $\geq 35$  kg/m<sup>2</sup> or 120% of the 95th percentile with clinically significant co-morbid conditions or BMI  $\geq 40$  kg/m<sup>2</sup> or 140% of the 95th percentile (whichever is lower) should be considered for MBS. Significant comorbidities include: obstructive sleep apnea (apnea-hypopnea index greater than 5), type 2 diabetes, idiopathic intracranial hypertension, nonalcoholic steatohepatitis (NASH), Blount's disease, slipped capital femoral epiphysis, GERD, or hypertension. The committee states a multidisciplinary team must also consider whether the patient and family have the ability and motivation to adhere to recommended treatments pre-and postoperatively, including consistent use of micronutrient supplements. The committee states adjustable gastric band and biliopancreatic diversion procedures in adolescents are less desirable choices based on the risk for reoperation. The adjustable gastric band is not currently FDA approved in patients under 18 years of age.

The short- and mid-term outcomes and complication rates of sleeve gastrectomy appear to be similar to those of other restrictive and malabsorptive procedures (Leyba, et al. 2011; Himpens, et al. 2010; D'Hondt, et al. 2011; Chouillard, et al. 2011). The procedure was proposed initially as the first step in a staged procedure for high-risk (super obese) patients, and longest follow-up data is available for these patients.

The ORBERA™ IntraGastric Balloon (Apollo Endosurgery, Inc.) received FDA approval in August 2015. The FDA approval is for patients with a BMI of 30-40 to assist in losing and maintaining weight. The ReShape Integrated Dual Balloon System (REShape medical, Inc.) also received FDA approval in July 2015. The device is also intended to facilitate weight loss in obese adult patients with a BMI of 30-40 who have been unsuccessful in losing weight through diet and exercise. Patients must also have one or more obesity related conditions such as diabetes, high blood pressure or high cholesterol. Both approved devices are considered temporary and should be removed after six months. Obalon Therapeutics received FDA approval for the Obalon Balloon System in September 2016. The Obalon Balloon System is a swallowable intraGastric balloon system indicated for temporary use to facilitate weight loss in adults with obesity (BMI of 30 – 40 kg/m<sup>2</sup>) who have failed to lose weight through diet and exercise. The System is intended to be used as an adjunct to a moderate intensity diet and behavior modification program. All balloons must be removed six months after the first balloon is placed. Peer-reviewed literature is currently insufficient to determine the efficacy and safety of an intraGastric balloon as a weight loss technique. A systematic review (Fernandes, et al. 2007) to assess the efficacy of the intraGastric balloon (IGB) found studies had very short follow-up and when compared to conventional management, IGB outcome data did not provide convincing evidence of a greater weight loss. More recent studies (e.g., REDUCE trial by

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 8 of 17**

Ponce, et al. 2015) found promising short-term (up to 48 weeks) results of weight loss when balloon insertion was compared to diet and exercise alone. In the Obalon clinical trial, 387 patients in the United States across fifteen clinical trial sites were randomized in a double-blind, sham-controlled study. The patients in the clinical trial received either three Obalon balloons or three sham placebo-like devices that looked similar to the balloons, but were filled with sugar. The patients in both groups were given minimal diet counseling of 25 minutes every three weeks. Both co-primary weight loss endpoints were met, with approximately 65% of patients who received the Obalon Balloon System experiencing clinically meaningful weight loss of at least 5% of their total body weight, which is twice as many people than in the sham-control group. Further studies are needed to demonstrate the long-term effects of utilizing intragastric balloon as a weight loss strategy.

Literature related to transoral gastroplasty for the treatment of obesity is limited; data is insufficient to provide conclusions on its safety and efficacy. The results of two studies (n=21, BMI range 35-53) were presented at the 2007 SAGES Annual Scientific Session. Follow-up for six patients at six months demonstrated an average weight loss of 31 pounds and an EWL of 24.9%. Sham-controlled trials are needed to further evaluate the utility of the EndoCinch device in obesity. Well-designed studies with long-term follow-up will be needed to measure the durability of the observed weight loss. Particularly, the stability of the gastric sutures remains unproven given the lack of long-term data.

There is not sufficient data published in the medical literature to draw conclusions about the safety and effectiveness of the mini-gastric bypass (also called loop gastric by-pass) procedure.

While preliminary data show that gastric plication has acceptable complication rates and weight loss outcomes in the short-term (e.g., Fried, et al. 2012; Skrekas, et al. 2011; Kourkoulos, et al. 2012; Talebpour, et al. 2012), additional well-designed comparative studies with established bariatric procedures are needed to determine its overall safety, efficacy and impact on health outcomes.

No controlled trials of single anastomosis duodeno-ileal bypass with sleeve gastrectomy have been identified. Some case series have reported on weight loss and other clinical outcomes up to five years post surgery. One larger series was published in 2015 (Sanchez-Pernaute, et al.) and reported on 97 patients with obesity and T2D. The authors reported that control of diabetes, defined as an HbA1c less than 6.0%, was achieved between 70% and 84% of patients at different time points. Remission rates were higher for patients on oral therapy than those on insulin, and were higher in patients with a shorter duration of diabetes. Currently, data is insufficient to provide conclusions on SADI-S safety and efficacy.

The AspireAssist obesity device (Aspire Bariatrics) received FDA approval in June 2016. Per the manufacturer, the AspireAssist device should not be used on patients with eating disorders, and it is not intended to be used for short durations in those who are moderately overweight. It is intended to assist in weight loss in patients aged 22 and older who are obese, with a body mass index of 35 to 55, and who have failed to achieve and maintain weight loss through non-surgical weight-loss therapy. The FDA reviewed results from a clinical trial of 111 patients treated with AspireAssist and appropriate lifestyle therapy, and 60 control patients who received only the lifestyle therapy. After one year, patients using AspireAssist lost an average of 12.1 percent of their total body weight compared to 3.6 percent for the control patients. There is insufficient literature to determine its overall benefit and safety as a bariatric procedure.

Transoral outlet reduction is being evaluated as an endoscopic revisional surgery in patients with weight regain following their primary bariatric procedure (e.g., gastric bypass). Preliminary results are promising with feasibility, safety and short-term efficacy being demonstrated in case series (Jirapino, et al. 2013; Thompson, et al. 2013, Kumar, et al. 2014). However, the long-term durability of the procedure still needs to be proven in larger studies.

The literature is insufficient to determine the long-term outcomes of stomach intestine pylorus sparing surgery (SIPS) in the treatment of morbid obesity.

Twelve months of abstinence is usually required for patients with a history of drug/alcohol abuse as liver toxins need to be avoided due to the higher rate of liver disease post bariatric surgery.

There is very little information available on the prevalence of asymptomatic liver disease in morbidly obese patients. The role of routine liver biopsy plays in patients undergoing bariatric surgery is not known. Liver biopsy is not routine practice in bariatric surgery. Impact on health outcomes has not been established through well-designed clinical trials. At this time, there is not sufficient evidence to support routine liver biopsy in patients undergoing bariatric surgery. Additional data from well-designed RCTs are needed to support the need for routine liver biopsy during bariatric surgical procedures.

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY****Policy Number: 7.01.29****Page: 9 of 17**

The British Society of Gastroenterology guidelines on the use of liver biopsy in clinical practice do not mention routine biopsy of the liver with bariatric surgery or indeed any other abdominal surgery. Recent American guidelines also do not endorse routine liver biopsies with abdominal surgery.

In a Choosing Wisely statement (2015), the American Society for Metabolic and Bariatric Surgery states the incidence of bile duct injury rates has increased since the introduction of laparoscopic cholecystectomy. The removal of normal and asymptomatic gallbladders at the time of bariatric surgery has not been shown to be necessary. In a 2016 retrospective cohort study, Sucandy et al. found a 22.7% incidence of subsequent biliary complications in patients who underwent biliopancreatic diversion and duodenal switch (BPD/DS).

**CODES**

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*
- *Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN)*

**CPT Codes**

<b>Code</b>	<b>Description</b>
43644	Laparoscopy, surgical, gastric restrictive procedure; with gastric bypass and Roux-en-Y gastroenterostomy (roux limb 150 cm or less)
43645	with gastric bypass and small intestine reconstruction to limit absorption
43770	Laparoscopy, surgical, gastric restrictive procedure; placement of adjustable gastric restrictive device (gastric band and subcutaneous port components)
43771	revision of adjustable gastric restrictive device component only
43772	removal of adjustable gastric restrictive device component only
43773	removal and replacement of adjustable gastric restrictive device-component only
43774	removal of adjustable gastric restrictive device and subcutaneous port components
43775	Laparoscopy, surgical, gastric restrictive procedure; longitudinal gastrectomy (ie, sleeve gastrectomy)
43842	Gastric restrictive procedure, without gastric bypass, for morbid obesity; vertical banded gastroplasty
43843	other than vertical –banded gastroplasty
43845	Gastric restrictive procedure with partial gastrectomy, pylorus-preserving duodenoileostomy and ileoileostomy (50 to 100 cm common channel) to limit absorption (biliopancreatic diversion with duodenal switch)
43846	Gastric restrictive procedure, with gastric bypass, for morbid obesity; with short limb (150 cm or less) Roux-en-Y gastroenterostomy
43847	with small bowel reconstruction to limit absorption
43848	Revision, open, of gastric restrictive procedure for morbid obesity, other than adjustable gastric restrictive device (separate procedure)
43886	Gastric restrictive procedure, open; revision of subcutaneous port components only
43887	removal of subcutaneous port component only
43888	removal and replacement of subcutaneous port component only

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY****Policy Number: 7.01.29****Page: 10 of 17**

Copyright © 2019 American Medical Association, Chicago, IL

**HCPCS Codes**

Code	Description
S2083	Adjustment of gastric band diameter via subcutaneous port by injection or aspiration of saline

**ICD10 Codes**

Code	Description
E66.01	Morbid (severe) obesity due to excess calories
E66.2	Morbid (severe) obesity with alveolar hypoventilation
Z68.35-Z68.45	Body mass index (BMI), 35.0-70 or greater, adult (code range)

**REFERENCES**

\*Abbatini F, et al. Long-term effects of laparoscopic sleeve gastrectomy, gastric bypass, and adjustable gastric banding on type 2 diabetes. Surg Endosc 2010 May;24(5):1005-10.

\*Abbatini F, et al. Type 2 diabetes in obese patients with body mass index of 30-35 kg/m<sup>2</sup>; sleeve gastrectomy versus medical treatment. Surg Obes Relat Dis 2012 Jun;8(3):20-4.

\*Abdelbaki TN, et al. Gastric plication for morbid obesity: a systematic review. Obes Surg 2012 Oct;22(10):1633-8.

Abu Dayyeh BK, et al. Endoscopic sleeve gastropasty alters gastric physiology and induces loss of body weight in obese individuals. Aliment Tract 2017;15:37-43.

\*Abu-Gazala S, et al. Conversion of failed gastric banding into four different bariatric procedures. Surg Obes Relat Dis 2011;doi:10.1016/j.soard.2011.01.009.

Adams TD, et al. Weight and metabolic outcomes 12 years after gastric bypass. N Engl J Med 2017 Sept 21;377(12):1143-1155.

Agency for Healthcare Research and Quality (AHRQ). Technology Assessment Program Project ID: OBST0816. Short- and long-term outcomes after bariatric surgery in the Medicare population. 2018 Sep 5. [https://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/ta/bariatric-surgery/bariatric-surgery-medicare-final-report.pdf] accessed 2/1/19.

\*Aikenhead A, et al. Effectiveness and cost-effectiveness of paediatric bariatric surgery: a systematic review. Clin Obes 2011 Feb;1(1):12-25.

\*Alami RS, et al. Is there a benefit in preoperative weight loss in gastric bypass patients? A prospective randomized trial. Surg Obes Relat Dis 2007 Mar-Apr;3(2):141-5.

Alfredo G, et al. Long-term multiple intragastric balloon treatment- a new strategy to treat morbid obese patients refusing surgery: prospective 6-year follow-up study. Surg Obes Relat Dis 2014 Mar-Apr;10(2):307-11.

Ali MR, et al. American Society for Metabolic and Bariatric Surgery position statement on intragastric balloon therapy endorsed by the Society of American Gastrointestinal and Endoscopic Surgeons. Surg Obes Relat Dis 2016 March-April;12(3):462-467.

Altieri MS, et al. Rate of revisions or conversion after bariatric surgery over 10 years in the state of New York. Surg Obes Relat Dis 2018 Apr;14(4):500-507.

American Society for Metabolic and Bariatric Surgery. Updated position statement on sleeve gastrectomy as a bariatric procedure. [http://asmbs.org/2012/06/sleeve-gastrectomy-as-a-bariatric-procedure-update/] assessed 1/28/19.

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 11 of 17**

\*American Society for Metabolic and Bariatric Surgery (ASMBS) Clinical issue Committee. ASMBS policy statement on gastric plication. Surg Obes Relat Dis 2011 May-Jun;7(3):262. updated 2015 Oct [<https://asmbs.org/resources/policy-statement-on-gastric-plication>] accessed 1/28/19.

American Society for Metabolic and Bariatric Surgery. ASMBS updated position statement on bariatric surgery in class I obesity. 2018 Aug. [<https://asmbs.org/resources/asmbs-updated-position-statement-on-bariatric-surgery-in-class-i-obesity>] accessed 1/28/19.

American Society for Metabolic and Bariatric Surgery. ASMBS pediatric metabolic and bariatric surgery guidelines. 2018 Aug. [<https://asmbs.org/resources/asmbs-pediatric-metabolic-and-bariatric-surgery-guidelines>] accessed 1/28/19.

Anderin C, et al. Weight loss before bariatric surgery and postoperative complications: data from the Scandinavian Obesity Registry (SOREg). Ann Surg 2015 May;261(5):909-13.

\*Angrisani L, et al. Laparoscopic adjustable gastric banding versus Roux-en-Y gastric bypass: 5-year results of a prospective randomized trial. Surg Obes Relat Dis 2007 Mar-Apr;3(2):127-32.

Barry RG, et al. Laparoscopic vertical sleeve gastrectomy: a 5-year veterans affairs review. Medicine 2017 Sept;96(35):e7508.

BlueCross BlueShield Association. Blue Distinction Centers of Bariatric Surgery. <https://www.bcbs.com/about-us/capabilities-initiatives/blue-distinction/blue-distinction-specialty-care#bariatric>] accessed 1/28/19.

BlueCross BlueShield Association. Bariatric surgery. Medical Policy Reference Manual. Policy #7.01.47. 2018 Feb 8.

\*BlueCross BlueShield Association Technology Evaluation Center (TEC). Newer techniques in bariatric surgery for morbid obesity: laparoscopic adjustable gastric banding, biliopancreatic diversion, and long-limb gastric bypass. 2005 Aug;20(5).

\*BlueCross BlueShield Association Technology Evaluation Center (TEC). Laparoscopic adjustable gastric banding in patients with body mass index less than 35 kg/m<sup>2</sup> with weight-related comorbidity. 2012 Oct;27(3).

\*BlueCross BlueShield Association Technology Evaluation Center (TEC). Bariatric surgery in patients with diabetes and body mass index less than 35 kg/m<sup>2</sup>. 2012 Oct;27(3).

\*Bohdjalian A, et al. Sleeve gastrectomy as sole and definitive bariatric procedure: 5-year results for weight loss and ghrelin. Obesity Surg 2010 May;20 (5):535-40.

Bower G, et al. Bariatric surgery and non-alcoholic fatty liver disease: a systematic review of liver biochemistry and histology. Obes Surg 2015;25(12):2280-2289.

\*Brethlauer S. ASMBS position statement on preoperative supervised weight loss requirements. Surg Obes Relat Dis 2011 May-Jun;7(3):257-60.

\*Brethlaauer S, et al. Transoral gastric volume reduction as intervention for weight management: 12-month follow-up of TRIM trial. Surg Obes Relat Dis 2012 Jun;8(3):296-304.

Carbajo MA, et al. Evaluation of weight loss indicators and laparoscopic one-anastomosis gastric bypass outcomes. Sci Rep 2018 Jan 31;8(1):1961.

\*Chakravarty PD, et al. Comparison of laparoscopic adjustable gastric banding (LAGB) with other bariatric procedures; a systematic review of the randomized controlled trials. Surgeon 2012 Jun;10(3):172-82.

Chang SH, et al. The effectiveness and risks of bariatric surgery: an updated systematic review and meta-analysis, 2003-2012. JAMA Surg 2014 Mar;149(3):275-87.

Changela K, et al. Peroral endoscopic reduction of dilated gastrojejunal anastomosis after bariatric surgery: techniques and efficacy. World J Gastrointest Endosc 2016 Feb 25;8(4):239-243.

Childerhose JE, et al. Adolescent bariatric surgery: a systematic review of recommendation documents. Surg Obes Relat Dis 2017 Oct;13(10):1768-1779.

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 12 of 17**

\*Choosing Wisely. American Society for Metabolic and Bariatric Surgery. Routine gallbladder removal. 2015 Jun 25. [<http://www.choosingwisely.org/clinician-lists/american-society-metabolic-bariatric-surgery-routine-gallbladder-removal/>] accessed 1/29/19.

\*Chouillard EK, et al. Laparoscopic Roux-en-Y gastric bypass versus laparoscopic sleeve gastrectomy for morbid obesity: case-control study. Surg Obes Relat Dis 2011 Jul-Aug;7(4):500-5.

Colquitt JL, et al. Surgery for weight loss in adults. Cochrane Database Syst Rev 2014 Aug 8;8:CD003641.

Cook FJ, et al. Long-term bone health after Roux-en-Y gastric bypass: a pilot study. Endocr Pract 2017 Sept;23(9):1077-1084.

Courcoulas AP, et al. Surgical vs medical treatments for type 2 diabetes mellitus: a randomized clinical trial. JAMA Surg 2014 Jul;149(7):707-15.

Courcoulas AP, et al. Three-year outcomes of bariatric surgery vs lifestyle intervention for type 2 diabetes mellitus treatment: a randomized clinical trial. JAMA Surg 2015 Oct;150(10):931-940.

Dardamanis D, et al. A retrospective comparative study of primary versus revisional Roux-en-Y gastric bypass: long-term results. Obes Surg 2018 Aug;28(8):2457-2464.

Della Penna A, et al. Ursodeoxycholic acid for 6 months after bariatric surgery is impacting gallstone associated morbidity in patients with preoperative asymptomatic gallstones. Obes Surg 2019 Jan 2. [Epub ahead of print]

\*D'Hondt M, et al. Laparoscopic sleeve gastrectomy as a single-stage procedure for the treatment of morbid obesity and the resulting quality of life, resolution of comorbidities, food tolerance, and 6-year weight loss. Surg Endosc 2011 Aug;25(8):2498-504.

Eid GM, et al. StomaphyX vs sham procedure for revisional surgery to reduce regained weight in Roux-en-Y gastric bypass patients: a randomized clinical trial. JAMA Surg 2014 Apr;149(4):372-9.

Ells LJ, et al. Surgery for the treatment of obesity in children and adolescents. Cochrane Database Syst Rev 2015 Jun 24;(6):CD011740.

Executive summary: Guidelines (2013) for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Obesity Society published by the Obesity Society and American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Based on a systematic review from the Obesity Expert Panel, 2013. Obesity 2014 Jul;22 Suppl 2:S5-39.

Fossell H, et al. A novel endoscopic weight loss therapy using gastric aspiration: results after 6 months. Endoscopy 2015 Jan;47(1):68-71.

\*Fried M, et al. Laparoscopic greater curvature plication (LGCP) for treatment of morbid obesity in a series of 244 patients. Obes Surg 2012 Aug;22(8):1298-307.

Garvey WT, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Comprehensive Clinical practice Guidelines for Medical Care of Patients with Obesity Executive Summary. Endocr Pract 2016 July;22(7):842-884.

Halperin F, et al. Roux-en-Y gastric bypass surgery or lifestyle with intensive medical management in patients with type 2 diabetes: feasibility and 1-year results of a randomized clinical trial. JAMA Surg 2014 Jul;149(7):716-26.

\*Hedberg J, et al. Superior weight loss and lower HbA1c 3 years after duodenal switch compared with Roux-en-Y gastric bypass-- a randomized controlled trial. Surg Obes Relat Dis 2012 May-Jun;8(3):338-43.

\*Himpens J, et al. Long-term results of laparoscopic sleeve gastrectomy for obesity. Ann Surg 2010 Aug;252(2):319-24.

\*Himpens J, et al. Long-term results of laparoscopic Roux-en-Y gastric bypass: evaluation after 9 years. Obes Surg 2012 Oct;22(10):1586-93.

Hsu CC, et al. Effect of bariatric surgery vs medical treatment on type 2 diabetes in patients with body mass index lower than 35: five-year outcomes. JAMA Surg 2015 Dec;150(12):1117-1124.

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 13 of 17**

Ibrahim AM, et al. Reoperation and Medicare expenditures after laparoscopic gastric band surgery. JAMA Surg 2017 Sept 1;152(9):835-842.

Inge TH, et al. Weight loss and health status 3 years after bariatric surgery in adolescents. N Engl J Med 2016 Jan 14;374(2):113-123.

Inge TH, et al. Comparison of surgical and medical therapy for Type 2 diabetes in severely obese adolescents. JAMA Pediatr 2018 May 1;172(5):452-460.

Jensen MD, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. Circulation 2014 Jun 24;129(25 Suppl 2):S102-38.

Ji Y, et al. A systematic review of gastric plication for the treatment of obesity. Surg Obes Relat Dis 2014 Nov-Dec;10(6):1226-1232.

\*Jirapinyo P, et al. Evaluation of an endoscopic suturing device for transoral outlet reduction in patients with weight regain following Roux-en-Y gastric bypass. Endoscopy 2013 Jul;45(7):532-6.

Jirapinyo P, et al. Gastrojejunal anastomotic reduction for weight regain in Roux-en-Y gastric bypass patients: physiological, behavioral, and anatomical effects of endoscopic suturing and sclerotherapy. Surg Obes Relat Dis 2016 Dec;12:1810-1816.

Juodeikis Z, et al. Long-term results after sleeve gastrectomy: a systematic review. Surg Obes Relat Dis 2017 Apr;13(4):693-699.

Kang JH, et al. Effectiveness of bariatric surgical procedures: a systematic review and network meta-analysis of randomized controlled trials. Medicine 2017 Nov;96(46):e8632.

Kashyap SR, et al. Metabolic effects of bariatric surgery in patients with moderate obesity and type 2 diabetes: analysis of a randomized controlled trial comparing surgery with intensive medical treatment. Diabetes Care 2013 Aug;36(8):2175-82.

\*Keating CL, et al. Cost-efficacy of surgically induced weight loss for the management of type 2 diabetes: a randomized controlled trial. Diabetes Care 2009 Apr;32(4):580-4.

Kim J, et al. American Society for Metabolic and Bariatric Surgery statement on single anastomosis duodenal switch. Surg Obes Relat Dis 2016 June; 12(5):944-945.

Kim SH, et al. Current status of intragastric balloon for obesity treatment. World J Gastroenterol 2016 June 28;22(24):5498-54504.

Kleiner DE, et al. Hepatic pathology among patients without known liver disease undergoing bariatric surgery: observations and a perspective from the longitudinal assessment of bariatric surgery (LABS) study. Semin Liver Dis 2014 Feb;34(1):98-107.

Koehestanie P, et al. The effect of the endoscopic duodenal-jejunal bypass liner on obesity and type 2 diabetes mellitus, a multicenter randomized controlled trial. Ann Surg 2014 Dec;260(6):984-992.

\*Kourkoulos M, et al. Laparoscopic gastric plication for the treatment of morbid obesity: a review. Minim Invasive Surg 2012;2012:696348.

\*Kumar N, et al. Comparison of a superficial suturing device with a full thickness suturing device for transoral outlet reduction. Gastrointest Endoscopy 2014 Jun;79(6):984-9.

Kumar N and Thompson CC. Transoral outlet reduction for therapy of weight regain after gastric bypass: long-term follow-up. Gastrointest Endosc 2015 April;83(4):776.

Kumar N, et al. Transoral outlet reduction for weight gain after gastric bypass: long-term follow-up. Gastrointest Endo 2016 April;83(4):776-779.

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 14 of 17**

Kumar N, et al. The role of endoscopic therapy in obesity management: intragastric balloons and aspiration therapy. Diabetes and metab Syndr Obes 2017 July 6;10:311-316.

\*Lee WJ, et al. Gastric bypass vs sleeve gastrectomy for type 2 diabetes mellitus: a randomized controlled trial. Arch Surg 2011 Feb;146(2):143-8.

\*Leyba JL, et al. Laparoscopic Roux-en-Y gastric bypass versus laparoscopic sleeve gastrectomy for the treatment of morbid obesity. A prospective study of 117 patients. Obes Surg 2011 Feb;21(2):212-6.

Lopez-Nava G, et al. Endoscopic sleeve gastroplasty for obesity: a multicenter study of 248 patients with 24 months follow-up. Obes Surg 2017 Oct;27(10):2649-2655.

\*Magouliotis DE, et al. Ursodeoxycholic acid in the prevention of gallstone formation after bariatric surgery: an updated systematic review and meta-analysis. Obes Surg 2017 Nov;27(11):3021–3030.

Mahawar KK, et al. Routine liver biopsy during bariatric surgery: an analysis of evidence base. Obes Surg 2016 Jan;26(1):177-181.

Manco M, et al. The benefit of sleeve gastrectomy in obese adolescents on nonalcoholic steatohepatitis and hepatic fibrosis. J Pediatr 2017 Jan;180:31-37.

Marin-Perez P, et al. Outcomes after laparoscopic conversion of failed adjustable gastric banding to sleeve gastrectomy or Roux-en-Y gastric bypass. Br J Surg 2014 Feb;101(3):254-60.

Martins Fernandes FA Jr, et al. Intragastric balloon for overweight patients. JLS 2016 Jan-Mar;20(1).

Maura D, et al. Effectiveness of intragastric balloon for obesity: a systematic review and meta-analysis based on randomized control trials. Surg Obes Relat Dis 2016 Feb;12(2):420-429.

Mingrone G, et al. Bariatric-metabolic surgery versus conventional medical treatment in obese patients with type 2 diabetes: 5-year follow-up of an open-label, single, centre, randomized controlled trial. Lancet 2016 Sept 5;386(9997):964-973.

Mitzman B, et al. Stomach intestinal pylorus sparing (SIPS) surgery for morbid obesity: retrospective analyses of our preliminary experience. Obes Surg 2016;1-7.

Morita S, et al. Prevalence of non-alcoholic fatty liver disease and steatohepatitis risk factors in patients undergoing bariatric surgery. Obes Surg 2015 Dec;25(12):2335-2343.

Moura D, et al. Effectiveness of intragastric balloon for obesity: a systematic review and meta-analysis based on randomized control trials. Surg Obes Relat Dis 2016 Feb;12(2):420-429.

Muller-Stich BP, et al. Surgical versus medical treatment of type 2 diabetes mellitus in nonseverely obese patients: a systematic review and meta-analysis. Ann Surg 2015 Mar;261(3):421-9.

Neichoy BT, et al. Stomach intestinal pylorus sparing surgery for morbid obesity. JLS 2018;22(1):e2017.

Noren E, et al. Aspiration therapy for obesity; a safe and effective treatment. BMC Obes 2016 Dec 28;3:56.

\*National Institutes of Health, National Heart, Lung and Blood Institute, Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Clinical Guidelines on the Identification, Evaluation and Treatment of Overweight and Obesity in Adults. Executive Summary. Bethesda, MD: National Institutes of Health; September 1998. [https://www.nhlbi.nih.gov/files/docs/guidelines/ob\\_gdlns.pdf](https://www.nhlbi.nih.gov/files/docs/guidelines/ob_gdlns.pdf) accessed 1/28/19.

National Institute for Health and Care Excellence. Implantation of a duodenal-jejunal bypass sleeve for managing obesity. IPG 471. Nov 2013 [<https://www.nice.org.uk/guidance/ipg471>] accessed 1/28/19.

National Institute for Health and Care Excellence. Single-anastomosis duodeno-ileal bypass with sleeve gastrectomy for treating morbid obesity. IPG 569. Nov 2016 [<https://www.nice.org.uk/guidance/ipg569/chapter/1-Recommendations>] accessed 1/28/19.

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 15 of 17**

Nedelcu M, et al. Laparoscopic sleeve gastrectomy: effect on long-term remission for morbidly obese patients with type 2 diabetes at 5-year follow-up. Surgery 2017 Oct;162(4):857-862.

Nobili V, et al. Indications and limitations of bariatric intervention in severely obese children and adolescents with and without nonalcoholic steatohepatitis: ESPGHAN hepatology Committee Position Statement. J Pediatr Gastroenterol Nutr 2015 April;60(4):550-561.

Novikov AA, et al. Endoscopic sleeve gastroplasty, laparoscopic sleeve gastrectomy, and laparoscopic band for weight loss: how do they compare? J Gastrointest Surg 2018 Feb;22(2):267-273.

Nystrom M, et al. Aspiration therapy as a tool to treat obesity: 1-to 4-year results in a 201-patient multicenter post-market European registry study. Obes Surg 2018 Jul;28(7):1860-1868.

Olbers T, et al. Laparoscopic Roux-en-Y gastric bypass in adolescents with severe obesity (AMOS): a prospective, 5-year, Swedish nationwide study. Lancet Diabetes Endocrinol 2017 March;5(3):174-183.

Osland E, et al. Weight loss outcomes in laparoscopic vertical sleeve gastrectomy (LVSG) versus laparoscopic Roux-en-Y gastric bypass (LRYGB) procedures: a meta-analysis and systematic review of randomized controlled trials. Surg Laparosc Endosc Percutan Tech 2017 Feb;27(1):8-18.

Palmisano S, et al. Intra-gastric balloon device: weight loss and satisfaction degree. Obes Surg 2016 Sept;26(9):2131-2137.

Patel L, et al. Outcomes following 50 consecutive endoscopic gastrojejunal revisions for weight gain following Roux-en-Y gastric bypass: a comparison of endoscopic suturing techniques for stoma reduction. Surg Endosc 2017 June;31(6):2667-2677.

Ponce J, et al. The REDUCE pivotal trial: a prospective, randomized controlled trial of a dual intra-gastric balloon for the treatment of obesity. Surg Obes Relat Dis 2015 Jul-Aug;11(4):874-81.

Puzziferri N, et al. Long-term follow-up after bariatric surgery: a systematic review. JAMA 2014 Sep 3;312(9):934-42.

Qi L, et al. Effects of bariatric surgery on glycemic and lipid metabolism, surgical complication and quality of life in adolescents with obesity: a systematic review and meta-analysis. Surg Obes Relat Dis 2017 Dec;13(12):2037-2055.

Qiu J, et al. Revisional bariatric surgery for weight regain and refractory complications in a single MBSAQIP accredited center: what are we dealing with? Obes Surg 2018 Sep;28(9):2789-2795.

Quezada N, et al. Safety and efficacy of the endoscopic duodenal-jejunal bypass liner prototype in severe or morbidly obese subjects implanted for up to 3 years. Surg Endosc 2018 Jan;32(1):260-267.

Raziel A, et al. Mid-term follow-up after laparoscopic sleeve gastrectomy in obese adolescents. Isr Med Assoc J 2014 Jan;16(1):37-41.

Risstad H, et al. Five-year outcomes after laparoscopic gastric bypass and laparoscopic duodenal switch in patients with body mass index of 50-60: a randomized clinical trial. JAMA Surg 2015 Apr;150(4):352-61.

Rohde U, et al. Effect of the Endobarrier gastrointestinal Liner on obesity and type 2 diabetes: a systematic review and meta-analysis. Diabetes Obes Metabol 2016 March;18(3):300-305.

Saber AA, et al. Efficacy of first-time intra-gastric balloon in weight loss: a systematic review and meta-analysis of randomized controlled trials. Obes Surg 2017 Feb;27(2):277-287.

Salminen P, et al. Effect of laparoscopic sleeve gastrectomy vs laparoscopic Roux-en-Y gastric bypass on weight loss at 5 years among patients with morbid obesity: the SLEEVEPASS randomized clinical trial. JAMA 2018 Jan 16;319(3):241-254.

Sanchez-Pernaute A, et al. Single-anastomosis duodenal bypass with sleeve gastrectomy (SADI-S) for obese diabetic patients. Surg Obes Relat Dis 2015 Sept-Oct;11(5):1092-1098.

Sharaiha RZ, et al. Endoscopic sleeve gastroplasty significantly reduces body mass index and metabolic complications in obese patients. Clin Gastroenterol Hepatol 2017 April;15(4):504-510.

**Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 16 of 17**

Schauer PR, et al. Bariatric surgery versus intensive medical therapy for diabetes-3 year outcomes. N Engl J Med 2014 May 22;370(21):2002-13.

Schauer PR, et al. Bariatric surgery vs. intensive medical therapy for long-term glycemic control and complications of diabetes. Final 5-year STAMPEDE trial results. American College of Cardiology 2016 Scientific Sessions; April 2, 2016; Chicago IL. Abstract 416-12.

Seki Y, et al. Prevalence of nonalcoholic steatohepatitis in Japanese patients with morbid obesity undergoing bariatric surgery. J Gastroenterol 2016 March;51(3):281-289.

Skogar ML, et al. Duodenal switch is superior to gastric bypass in patients with super obesity when evaluated with the bariatric analysis and reporting outcome system (BAROS). Obes Surg 2017 Sept;27(9):2308-2316.

\*Skrekas G, et al. Laparoscopic gastric greater curvature plication: results and complications in a series of 135 patients. Obes Surg 2011 Nov;21(11):1657-63.

Sogg S, et al. Recommendations for the presurgical psychosocial evaluation of bariatric surgery patients. ASMBS Guidelines/Statements. Surg Obes Relat Dis 2016 Dec;12:731-749.

\*Solomon H, et al. Benefits to patients choosing preoperative weight loss in gastric bypass surgery: new results of a randomized trial. J Am Coll Surg 2009 Feb;208(2):241-5.

Souto KP, et al. Nonalcoholic fatty liver disease in patients with different baseline glucose status undergoing bariatric surgery: analysis of intraoperative liver biopsies and literature review. Surg Obes Relat Dis 2018 Jan;14(1):66-73.

Stroh C, et al. Results of more than 11,800 sleeve gastrectomies: data analysis of the German Bariatric Surgery registry. Ann Surg 2016 May;263(5):949-955.

\*Sucandy I, et al. Risk of biliary events after selective cholecystectomy during biliopancreatic diversion with duodenal switch. Obes Surg 2016 Mar;26(3):531-7.

Sullivan S, et al. Randomized sham-controlled trial evaluating efficacy and safety of endoscopic gastric plication for primary obesity: The ESSENTIAL trial. Obesity 2017 Feb;25(2):294-301.

Switzer NJ, et al. Sleeve gastrectomy and Type 2 diabetes mellitus: a systematic review of long-term outcomes. Obes Surg 2016 July;26(7):1616-1621.

\*Talebpour M, et al. Twelve year experience of laparoscopic gastric plication in morbid obesity: development of the technique and patient outcomes. Ann Surg Innov Res 2012;Aug 22;6(1):7.

Talebpour M, et al. Comparison of short-term effectiveness and postoperative complications: laparoscopic gastric plication vs laparoscopic sleeve gastrectomy. Obes Surg 2018 Apr;28(4):996-1001.

\*Tarnoff M, et al. Open label, prospective, randomized controlled trial of an endoscopic duodenal-jejunal bypass sleeve versus low calorie diet for pre-operative weight loss in bariatric surgery. Surg Endosc 2009 Mar;23(3):650-6.

Tate CM, et al. Intra-gastric balloon treatment for obesity: review of recent studies. Adv Ther 2017 Aug;34(8):1859-1875.

Thereaux J, et al. Long-term follow-up after bariatric surgery in a national cohort. Br J Surg 2017 Sept;104(10):1362-1371.

\*Thompson CC, et al. Endoscopic suturing for transoral outlet reduction increases weight loss after Roux-en-Y gastric bypass surgery. Gastroenterology 2013 Jul;145(1):129-37.

Thompson CC, et al. Percutaneous gastrostomy device for the treatment of Class II and Class III obesity: results of a randomized controlled trial. Am J Gastroenterol 2017 March;112(3):447-457.

\*Tsilrine VB, et al. How frequently and when do patients undergo cholecystectomy after bariatric surgery? Surg Obes Relat Dis 2014 Mar-Apr;10(2):313-21.

Tustumi F, et al. Cholecystectomy in patients submitted to bariatric procedure: a systematic review and meta-analysis. Obes Surg 2018 Oct;28(10):3312-3320.

## **Medical Policy: SURGICAL MANAGEMENT OF OBESITY**

**Policy Number: 7.01.29**

**Page: 17 of 17**

Van Rutte PW, et al. Outcomes of sleeve gastrectomy as a primary bariatric procedure. Br J Surg 2014 May;101(6):661-8.

Wang MC, et al. Laparoscopic Roux-en-Y gastric bypass versus sleeve gastrectomy for obese patients with Type 2 diabetes: a meta-analysis of randomized controlled trials. Am Surg 2015 Feb;81(2):166-71.

Wang GF, et al. Predictive factors of type 2 diabetes mellitus remission following bariatric surgery: a meta-analysis. Obes Surg 2015 Feb;25(2):199-208.

Wu GZ, et al. Meta-analysis of bariatric surgery versus non-surgical treatment for type 2 diabetes mellitus. Oncotarget 2016 Dec 27;7(52):87511-87522.

Yan Y, et al. Roux-en-Y gastric bypass versus medical treatment for type 2 diabetes mellitus in obese patients: a systematic review and meta-analysis of randomized controlled trials. Medicine 2016 April;95(17):e3462.

Yorke E, et al. Intra-gastric balloon for management of severe obesity: a systematic review. Obes Surg 2016 Sept;26(9):2248-2254.

Young MT, et al. Use and outcomes of laparoscopic sleeve gastrectomy vs laparoscopic gastric bypass: analysis of the American College of Surgeons NSIP. J Am Coll Surg 2015 May;220(5):880-5.

Zellmer JD, et al. Is laparoscopic sleeve gastrectomy a lower risk bariatric procedure compared with laparoscopic Roux-en-Y gastric bypass? A meta-analysis. Am J Surg 2014 Dec;208(6):903-10.

Zheng Y, et al. Short-term effects of intra-gastric balloon in association with conservative therapy on weight loss: a meta-analysis. J Transl Med 2015 July 29;13:246.

\*Key Article

### **KEY WORDS**

Adjustable gastric band, Aspiration therapy, Bariatric, Endobarrier, Gastric bypass, Gastric plication, Imbrication, intra-gastric balloon, Lap Band, ROSE, Roux-en-Y, SIPS, Sleeve gastrectomy, Transoral outlet reduction.

### **CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a National Coverage Determination (NCD) for bariatric surgery. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=57&ncdver=5&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&CptHcpcsCode=36514&bc=gAAAABAAAAAAAA%3d%3d&>

There is also an article from the National Government Services contractor related to sleeve gastrectomy. Please refer to the following web site:

[https://www.cms.gov/medicare-coverage-database/details/article-details.aspx?articleId=52447&ver=12&Ctrctr=299&ContrVer=1&CtrctrSelected=299\\*1&name=National+Government+Services%2c+Inc.+\(National+Government+Services%2c+Inc.+\(13102%2c+A+and+B+and+HHH+MAC%2c+J+-+K\)\)&LCtrctr=299\\*1&DocType=Active&bc=AgAAAIAgAAA&](https://www.cms.gov/medicare-coverage-database/details/article-details.aspx?articleId=52447&ver=12&Ctrctr=299&ContrVer=1&CtrctrSelected=299*1&name=National+Government+Services%2c+Inc.+(National+Government+Services%2c+Inc.+(13102%2c+A+and+B+and+HHH+MAC%2c+J+-+K))&LCtrctr=299*1&DocType=Active&bc=AgAAAIAgAAA&)