

MEDICAL POLICY

SUBJECT: CONTINUOUS PASSIVE MOTION DEVICE IN THE HOME SETTING	EFFECTIVE DATE: 10/18/01 REVISED DATE: 11/21/02, 09/18/03, 06/17/04, 04/21/05, 06/22/06, 08/23/07, 10/23/08, 08/27/09, 08/26/10, 08/25/11, 10/25/12, 08/22/13, 08/28/14, 08/27/15, 10/27/16, 10/26/17 ARCHIVED DATE: 08/23/18
POLICY NUMBER: 1.01.02 CATEGORY: Equipment/Supplies	PAGE: 1 OF: 3
<ul style="list-style-type: none">• <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i>• <i>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</i>• <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i>	

POLICY STATEMENT:

- I. Based upon our criteria and assessment of peer-reviewed literature a Continuous Passive Motion (CPM) device is **medically appropriate** in the home setting following surgery under conditions of low postoperative mobility or inability to comply with rehabilitation exercises or during the non-weight bearing rehabilitation period for the following:
 - A. Total knee arthroplasty (TKA) or equivalent open knee surgery;
 - B. Anterior cruciate ligament (ACL) reconstruction;
 - C. Open reduction and internal fixation (ORIF) of tibial plateau or distal femur fractures involving the knee joint;
or
 - D. Surgical release of arthrofibrosis/adhesive capsulitis or manipulation under anesthesia of the knee until the member is participating in an active physical therapy (PT) program.
- II. Use of the CPM device beyond 21 days post-operatively or 21 days after manipulation under anesthesia is not supported by the medical literature, as it does not improve patient outcomes, and is considered **not medically necessary**.
- III. CPM as an adjunct to conventional physical therapy for any other indication (e.g., rotator cuff, metacarpophalangeal or temporomandibular joint) has not been medically proven to be effective and is considered **not medically necessary**.

POLICY GUIDELINES:

- I. Contractual coverage of durable medical equipment is required.
- II. The CPM device is only allowable for up to 21 days if the patient is compliant with the regimen.

DESCRIPTION:

The restoration of joint range of motion following surgery or trauma is dependent upon rehabilitation. Delay in rehabilitation may subsequently result in poor joint function or immobility. Passive motion, a treatment component of joint rehabilitation, may be performed by a physical therapist or accomplished with a continuous passive motion (CPM) device. The device moves the joint (e.g., flexion, extension) without patient assistance, continuously. The device is held in place across the affected joint by Velcro straps. An electrical power unit is used to set the variable range of motion (ROM) and speed.

To some extent, CPM devices and physical therapists are interchangeable means of delivering passive motion. The preference for one or the other mode of delivery may be determined by considerations related to the organization of services or resource allocation (e.g., staffing, timing of discharge, patient access to physical therapy). Such use of a continuous passive motion device as a substitute for a physical therapist for delivering passive motion should be distinguished from use of the device as an adjunct to physical therapy, in which the objective is to increase the duration and intensity of passive motion in order to achieve outcomes superior to that achieved by conventional physical therapy programs.

A variety of CPM devices are available.

<p>SUBJECT: CONTINUOUS PASSIVE MOTION DEVICE IN THE HOME SETTING</p> <p>POLICY NUMBER: 1.01.02 CATEGORY: Equipment/Supplies</p>	<p>EFFECTIVE DATE: 10/18/01 REVISED DATE: 11/21/02, 09/18/03, 06/17/04, 04/21/05, 06/22/06, 08/23/07, 10/23/08, 08/27/09, 08/26/10, 08/25/11, 10/25/12, 08/22/13, 08/28/14, 08/27/15, 10/27/16, 10/26/17 ARCHIVED DATE: 08/23/18</p> <p>PAGE: 2 OF 3</p>
---	--

RATIONALE:

There is controversy in the published studies on the use of the CPM device. While some studies demonstrate no benefit beyond that of conventional physical therapy, others find that use of CPM is effective. There is evidence in the published studies that CPM as an adjunct to standard physical therapy used immediately following TKA, anterior cruciate ligament (ACL) reconstruction, and open reduction and internal fixation (ORIF) of tibial plateau or distal femur fractures involving the knee joint does improve net health outcomes beyond the benefit of physical therapy alone. Studies of postoperative use of CPM for other procedures (e.g., rotator cuff repair, metacarpophalangeal joint arthroplasty) and for stroke rehabilitation do not permit conclusions that the CPM device is effective.

CODES: Number Description

Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.

CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.

Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT: No specific code(s)

Copyright © 2018 American Medical Association, Chicago, IL

HCPCS: E0935 Continuous passive motion exercise device for use on the knee only
E0936 (NMN) Continuous passive motion exercise device for use other than knee

ICD10: M17.0-M17.9 Osteoarthritis of knee (code range)
M23.50 Chronic instability of knee, unspecified knee

REFERENCES:

BlueCross BlueShield Association. Continuous Passive Motion (CPM) in the home setting. Medical Policy Reference Manual Policy #1.01.10. 2018 Mar 08.

Boese CK, et al. The efficacy of continuous passive motion after total knee arthroplasty: a comparison of three protocols. J Arthroplasty 2014;29(6):1158-62.

Brennan GP, et al. Outpatient rehabilitation care process factors and clinical outcomes among patients discharged home following unilateral total knee arthroplasty. J Arthroplasty 2015 May;30(5):885-90.

Chaudry H, and Bhandari M. Cochrane in CORR (®): Continuous passive motion following total knee arthroplasty in people with arthritis (Review). Clin Orthop Relat Res. 2015 Nov;473(11):3348-54.

De Roo PJ, et al. Passive mobilization after arthroscopic rotator cuff repair is not detrimental in the early postoperative period. Acta Orthop Belg. 2015 Sep;81(3):485-92.

*Du Plessis M , et al. The effectiveness of continuous passive motion on range of motion, pain and muscle strength following rotator cuff repair: a systematic review. Clin Rehabil 2011 Apr;25(4):291-302.

Ekim AA, et al. Continuous passive motion in adhesive capsulitis patients with diabetes mellitus: A randomized controlled trial. J Back Musculoskelet Rehabil. 2016 Nov 21;29(4):779-786.

*Garafalo R, et al. Effects of one month continuous passive motion after arthroscopic rotator cuff repair: results at one year follow-up of a prospective randomized study. Musculoskel Surg 2010;94(Suppl 1):S79–83.

<p>SUBJECT: CONTINUOUS PASSIVE MOTION DEVICE IN THE HOME SETTING</p> <p>POLICY NUMBER: 1.01.02 CATEGORY: Equipment/Supplies</p>	<p>EFFECTIVE DATE: 10/18/01 REVISED DATE: 11/21/02, 09/18/03, 06/17/04, 04/21/05, 06/22/06, 08/23/07, 10/23/08, 08/27/09, 08/26/10, 08/25/11, 10/25/12, 08/22/13, 08/28/14, 08/27/15, 10/27/16, 10/26/17 ARCHIVED DATE: 08/23/18</p> <p>PAGE: 3 OF: 3</p>
---	---

Harvey La, et al. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev* 2014;(3):CD004260.

He ML, et al. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty. *Cochrane Database Syst Rev* 2014;7:CD008207.

Herbold JA, et al. Randomized controlled trial of the effectiveness of continuous passive motion after total knee replacement. *Arch Phys Med Rehabil* 2014 Jul;95(7):1240-5.

Herbold JA, et al. Effectiveness of continuous passive motion in an inpatient rehabilitation hospital after total knee replacement: a matched cohort study. *PM R* 2012 Oct;4(10):719-25.

*Koulalis D, et al. Autologous chondrocyte transplantation for osteochondritis dissecans of the talus. *Clin Orthop* 2002 Feb;(395):186-92.

*Lenssen AF, et al. Continuous passive motion following primary total knee arthroplasty: short-and long-term effects on range of motion. *Phys Ther Rev* 2003;8:113-21.

*Maloney GE, et al. Effect of a passive jaw motion device on pain and range of motion in TMD patients not responding to flat plane intraoral appliances. *Cranio* 2002 Jan;20(1):55-66.

Maniar RN, et al. To use or not to use continuous passive motion post-total knee arthroplasty presenting functional assessment results in early recovery. *J Arthroplasty* 2012 Feb;27(2):193-200.e1.

*Milne S, et al. Continuous passive motion following total knee arthroplasty. *Cochrane Database of Syst Rev* 2003;(2):CD004260.

Mistry JB, et al. Rehabilitative guidelines after total knee arthroplasty: a review. *J Knee Surg.* 2016 Apr;29(3):201-17.

*Nuyens G, et al. Reduction of spastic hypertonia during repeated passive knee movements in stroke patients. *Arch Phys Med Rehabil* 2002 Jul;83(7):930-5.

Riboh JC, et al. Early passive motion versus immobilization after arthroscopic rotator cuff repair. *Arthroscopy* 2014 Aug;30(8):997-1005.

Ross D, et al. Rehabilitation following arthroscopic rotator cuff repair: a review of current literature. *J Am Acad Orthop Surg* 2014 Jan;22(1):1-9.

Yi A, et al. A comparison of rehabilitation methods after arthroscopic rotator cuff repair: a systematic review. *Sports Health.* 2015 Jul;7(4):326-34.

* key article

KEY WORDS: Continuous passive motion, CPM, Knee.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently no National Coverage Determination (NCD) or Local Coverage Determination (LCD) for Continuous Passive Motion Devices. Please refer to the following NCD for Durable Medical Equipment Reference List website for Medicare Members: [http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=190&ncdver=2&NCAId=8&ver=7&NcaName=Augmentative+and+Alternative+Communication+\(AAC\)+Devices+for+Speech+Impairment&bc=ACAAAAAIAAA&](http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=190&ncdver=2&NCAId=8&ver=7&NcaName=Augmentative+and+Alternative+Communication+(AAC)+Devices+for+Speech+Impairment&bc=ACAAAAAIAAA&)