

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	LOW-DOSE COMPUTED TOMOGRAPHY (LDCT) FOR LUNG CANCER SCREENING
Policy Number	6.01.19
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	06/20/02, 05/21/03, 05/19/04, 05/18/05, 03/16/06, 12/21/06, 08/16/07, 06/19/08, 06/18/09, 12/20/12, 10/17/13, 06/19/14, 06/18/15
Archived Date	02/18/16
Edited Date	01/19/17, 01/18/18, 01/17/19
Deleted	05/27/10 – 12/20/12
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and review of the peer-reviewed literature, annual lung cancer screening using Low Dose CT is considered **medically appropriate** for high risk individuals who have not received a Low Dose CT lung screening in the past 12 months and have no signs or symptoms suggestive of underlying lung cancer, and are able and willing to undergo curative lung surgery. High risk individuals are defined as between age 55-80 years with 30 pack year history of smoking cigarettes who are either:
 1. a current smoker; or
 2. have quit smoking within the past 15 years.
- II. Based upon our criteria and review of the peer-reviewed literature, computer aided detection (CAD) has not been medically proven to be effective to improve the accuracy of CT scanning in screening for lung cancer and therefore is considered **investigational**.

Refer to Corporate Medical Policy #6.01.13, Computed Tomography (EBCT, Spiral CT, MDCT) to Detect Coronary Artery Calcification (Coronary Calcium Scoring).

POLICY GUIDELINES

- I. Lung cancer screening using Low Dose CT should be discontinued when:
 - A. An individual has not smoked for more than 15 years; or
 - B. An individual develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.
- II. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Spiral (helical or low dose) computed tomography (CT) is used to diagnose lung cancer in symptomatic persons. Published data has indicated that a low radiation dose spiral CT is capable of detecting abnormalities, including those suggestive of lung cancer, in asymptomatic, high-risk individuals. Electron beam CT (EBCT) is also used, however, spiral CT is the modality most commonly reported in the literature.

Spiral CT has several technical advantages over conventional CT, which enhances its clinical role. Imaging can be performed during a 20-second breath hold. The X-ray tube rotates continuously around the patient while the table and patient slowly move through the scanner. Each rotation takes 0.7-1 second therefore large volumes can be covered during a single scan. A volume data set is obtained as the tube-detector system traces a helical or spiral path. If the scan is performed during a single breath hold, it will be virtually free of misregistration artifacts. Radiation exposure is comparable to that absorbed during mammography. The entire examination is performed in less than 10 minutes.

The outcomes proposed for measuring the efficacy of spiral CT are:

- I. Detection of smaller and presumably earlier stage tumors than chest x-ray,
- II. Reduction in mortality of lung cancer patients, exposure to a lower dose of radiation than high resolution CT, no intravenous contrast needed, and
- III. Less expensive than high resolution CT and minimally more expensive than chest x-ray.

RATIONALE

Spiral CT systems such as the LightSpeed Plus CT System (General Electric Medical Systems) are approved by the U.S. Food and Drug Administration (FDA). In February 2004, the FDA approved the R2 Technology ImageChecker CT software system as a technique to assist in the detection of lung nodules on multi-detector CT scans of the chest.

While spiral computed tomography is proposed as an alternative to chest x-ray and high resolution CT in the screening of lung cancer, published clinical trials do not provide evidence to support the efficacy of spiral CT over conventional screening methods in reducing mortality from lung cancer. Studies support that spiral CT scanning is more sensitive than chest x-rays in identifying lung lesions. However, there is inadequate data to indicate whether early identification of lung cancer will lead to decreased cancer mortality. Because no lung cancer mortality data exists for spiral CT, there is insufficient evidence to advocate mass screening with spiral CT for individuals at elevated risk of lung cancer.

To be a valuable screening tool, it is not sufficient that spiral CT accurately detect malignant pulmonary nodules at an earlier stage. The technology must also demonstrate potential to prolong lung cancer survival time and to reduce disease-associated mortality. At this time, the available evidence from clinical research trials does not indicate clinically significant benefits or cost-effectiveness associated with spiral CT detection of lung cancer by routine screening.

Results from the International Early Lung Cancer Action Project (ELCAP) reported that of 31,657 asymptomatic patients who underwent a baseline and then annual CT scan for detection of lung cancer, a diagnosis of lung cancer was found in 484 patients (1.53% of the study population). Participants were enrolled at multiple sites worldwide, including the U.S., Japan and Europe. The study did not use a comparison group, such as screening with chest x-ray, to clearly demonstrate that there is any benefit from annual CT exams, and was non-randomized. Risks involved with CT screening are increased radiation exposure and the extent of evaluation of false-positive scans including needle biopsy and surgery to further evaluate a positive finding. Randomized controlled trials are needed to determine if use of this procedure improves survival and to assess the overall impact of various alternatives. One such trial, the National Lung Screening Trial, is in progress.

The National Lung Screening Trial (NLST) was a large well-conducted trial comprising a total of 53,454 current or former smokers from 33 sites in the United States had been randomly assigned to screening in 3 consecutive years with either a chest x-ray or low-dose spiral CT. Study eligibility included age between 55 and 74 years, a history of cigarette smoking of at least 30 pack-years and, for former smokers, quitting within the past 15 years. Individuals with a previous diagnosis of lung cancer or who had signs and/or symptoms suggestive of lung cancer were excluded. The trial results found a statistically significantly lower rate of lung cancer mortality with 3 annual CT screens compared to chest radiographs; the number needed to screen (NNS) to prevent one lung cancer death was 320 (95% CI: 193 to 934). The study also found a statistically significant but modestly lower overall mortality in low-dose CT group. There was a high rate of follow-up imaging tests but relatively low rates of invasive tests. There were few major complications reported after invasive testing, although major complications that did occur were not well-characterized. The rates of other potential complications, in particular radiation-induced cancers, are not yet known. Findings of the trial cannot be generalized to other populations, e.g., younger individuals or lighter smokers. The NLST evaluated the utility of a series of 3 annual CT screens; the efficacy of other screening regimens is not known.

Medical Policy: LOW-DOSE COMPUTED TOMOGRAPHY (LDCT) FOR LUNG CANCER SCREENING Policy Number: 6.01.19

Page: 3 of 6

Danish Lung Cancer Screening Trial randomized a total of 4,104 current or former smokers to screening with annual low-dose CT for 5 years or no screening during 2004 and 2006; lung cancer mortality was the primary outcome measure. Among the 2,052 individuals who received baseline CT scans, 179 (8.7%) had positive findings; a large proportion of these findings (162 of 179, 91%) were false positive. Seventeen individuals (0.8%) were found to have lung cancer; 10 cases were stage I disease.

On December 31, 2013, the U.S. Preventive Services Task Force (USPSTF) published their recommendation statement on lung cancer screening. The Task Force recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery. (B recommendation). The USPSTF concluded with moderate certainty that annual screening for lung cancer with LDCT is of moderate net benefit in asymptomatic persons who are at high risk for lung cancer based on age, total cumulative exposure to tobacco smoke, and years since quitting smoking. The moderate net benefit of screening depends on limiting screening to persons who are at high risk, the accuracy of image interpretation being similar to that found in the NLST (National Lung Screening Trial), and the resolution of most false-positive results without invasive procedures.

The National Comprehensive Cancer Network Lung Cancer Screening Guideline (2014) recommends that high risk individuals should be screened; however moderate- and low-risk should not currently be screened. High risk individuals are defined as between age 55-74 years with a 30 pack year history of smoking tobacco and if former smoker, have quit within 15 years. This is a category I recommendation because individuals are selected based on the NLST inclusion criteria. Individuals age 50 years or older, 20 or more pack year history of smoking tobacco and one additional risk factors. Risk factors include exposure to high radon, occupational exposure to either silica, cadmium, asbestos, arsenic, beryllium, chromium, diesel fumes or nickel, personal history of lung cancer, lymphoma, head and neck cancer, family history of lung cancer or personal history of COPD or pulmonary fibrosis. This is a category 2B recommendation because these individuals are selected based on lower level evidence and some panel members would not recommend LDCT for these individuals.

Professional organizations. A collaborative initiative of the American Cancer Society (ACS), the American College of Chest Physicians (ACCP), the American Society of Clinical Oncology (ASCO) developed the following recommendations: For smokers and former smokers aged 55 to 74 years who have smoked for 30 pack-years or more and either continue to smoke or have quit within the past 15 years, annual screening with low-dose computed tomography (LDCT) are suggested to be offered over both annual screening with chest radiograph or no screening, but only in settings that can deliver the comprehensive care provided to National Lung Screening Trial (NLST) participants. (Grade of recommendation: 2B.) For individuals who have accumulated fewer than 30 pack years of smoking or are either younger than 55 years or older than 74 years, or individuals who quit smoking more than 15 years ago, and for individuals with severe comorbidities that would preclude potentially curative treatment, limit life expectancy, or both, CT screening is suggested to not be performed. (Grade of recommendation: 2C.)

The Lung Association (2012) recommends lung cancer screening with low-dose CT scans for people who meet the following criteria: current or former smokers (aged 55 to 74 years), with a smoking history of at least 30 pack-years (that is, an average of a pack a day for 30 years) and with no history of lung cancer. The Lung Association emphasizes that only CT scans are recommended and that chest X-rays should not be used for lung cancer screening. The Lung Association recognizes that while low dose CT scans may save lives, screening for lung cancer should not be recommended for everyone as many known and unknown risks may be associated with the screening and subsequent medical evaluation.

The American Cancer Society Interim Guidance on Lung Cancer Screening(2012) recommend that adults between the ages of 55-74 *who meet the eligibility criteria of the NLST* and are concerned about their risk of lung cancer may consider screening for early lung cancer detection. With their physician or other primary provider, individuals interested in screening should weigh the currently known benefits of LDCT screening with the currently known limitations and risks and make a shared decision as to whether they should be screened for lung cancer.

The American Association for Thoracic Surgery guidelines for cancer screening using low-dose computed tomography scans for lung cancer survivors and other high risk groups (2012) recommends annual lung cancer screening to begin at

Medical Policy: LOW-DOSE COMPUTED TOMOGRAPHY (LDCT) FOR LUNG CANCER SCREENING Policy Number: 6.01.19

Page: 4 of 6

age 55 years for smokers and former smokers with a 20 pack-year history of smoking which may continue to age 79 years (Level 1 evidence). Annual lung cancer low-dose CT should be performed in patients who have been treated for a primary bronchogenic carcinoma without recurrence 4 years post- radiographic surveillance or in patients aged 50-79 years with a 20 pack-year smoking history and other factors (e.g., COPD with FEV1 of 70% or less than predicted, environmental and occupational exposures, prior cancer or thoracic radiation, or genetic or family history) that produce a cumulative risk of developing lung cancer that is 5% or more over the following 5 years (Level 2 evidence).

Computer Aided Detection. The use of computer-aided detection (CAD) software may assist in lung cancer screening. However, there is insufficient evidence to determine whether CAD technology may improve the accuracy of CT scanning interpretation. While CAD systems have been shown to detect additional lung nodules compared to the results of human readers alone, the issue is how many detected nodules are lung cancers. The effectiveness of CAD in detecting lung cancer has not been fully investigated. High-quality randomized trials examining the effect of CAD systems for CT scans on lung cancer morbidity and mortality are necessary to determine the true impact of this technology on health outcomes.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
71250	Computerized tomography, thorax; without contrast material (*NMN with V16.1, Z80.1)

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
G0296	Counseling visit to discuss need for lung cancer screening (LDCT) using low dose CT scan (service is for eligibility determination and shared decision making)
G0297	Low dose CT scan (LDCT) for lung cancer screening

ICD10 Codes

Code	Description
Z12.2	Special screening for malignant neoplasms, respiratory organs
Z80.1	Family history of malignant neoplasm; trachea, bronchus and lung

REFERENCES

Previously titled Spiral Computed Tomography (CT) for Lung Cancer Screening.

Aberle DR, et al. Reduced lung cancer mortality with low-dose computed tomographic screening. N Engl J Med 2011;365:395-409.

Aberle DR, et al. Results of the two incidence screenings in the National Lung Screening Trial. N Engl J Med 2013;369(10):920-31.

*Alberts WM, et al. Diagnosis and management of lung cancer executive summary: ACCP evidence-based clinical practice guidelines (2nd edition). Chest 2007 Sep;132(3 Suppl):1S-19S.

Ashraf H., et al. Smoking habits in the randomized Danish Lung Cancer Screening Trail with low-dose CT: final results after a 5-year screening programme. Thorax 2014 Jun;69(6):574-9.

Medical Policy: LOW-DOSE COMPUTED TOMOGRAPHY (LDCT) FOR LUNG CANCER SCREENING Policy Number: 6.01.19

Page: 5 of 6

Bach PB, et al. Benefits and harms of CT screening for lung cancer a systematic review. JAMA. 2012 Jun 13;307(22):2418-29.

Blue Cross BlueShield Association. Screening for lung cancer using CT scanning or chest radiographs. Medical Policy Reference Manual. Policy #6.01.30. Archived 2015 Nov 12.

CancerNet. Randomized study comparing annual chest x-rays and annual spiral CT scans in patients at high risk for lung cancer. Protocol ID: JHL-45199, NCI-V00-1600.

Christe A, et al. Lung cancer screening with CT: evaluation of radiologists and different computer assisted detection software (CAD) as first and second readers for lung nodule detection at different dose levels. Eur J Radiol 2013 Dec;82(12):e873-8.

*Clark LP, et al. National Cancer Institute initiative: lung image database resource for imaging research. Acad Radiol 2001 May;8(5):447-50.

Detterbeck FC, et al. Screening for lung cancer: diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013 May;143(5 Suppl):e78S-e92S.

*Diederich S, et al. Screening for early lung cancer with low-dose spiral CT: prevalence in 817 asymptomatic smokers. Radiol 2002 Mar;222(3):773-81.

Field JK, et al. CT screening for lung cancer: countdown to implementation. Lancet Oncol 2013; 14(13):e591-e600.

Freidrich MJ. Task force recommends targeted lung cancer screening. JAMA 2013 Sep 4;310(9):892-3.

Fu C, Liu Z, Zhu F, et al. A meta-analysis: is low-dose computed tomography a superior method for risky lung cancers screening population? Clin Respir J 2016 May;10(3):333-41.

Goulart BH, et al. Moving beyond the national lung screening trial: discussing strategies for implementation of lung cancer screening programs. Oncologist 2013;18(8):941-6.

*International Early Lung Cancer Action Program Investigators; Henschke CI, et al. Survival of patients with stage I lung cancer detected on CT screening. NEJM 2006 Oct;355(17):1763-71.

Jaklitsch MT, et al. The American Association for Thoracic Surgery guidelines for lung cancer screening using low-dose computed tomography scans for lung cancer survivors and other high-risk groups. J Thorac Cardiovasc Surg 2012;144(1):33-8.

Kanne JP. Screening for lung cancer: what have we learned? AJR Am J Roentgenol 2014 Mar;202(3):530-5.

Manser R, et al. Screening for lung cancer. Cochrane Database Syst Rev 2013 Jun 21;6:CD001991.

Mazzone PJ, et al. Lung cancer screening with computer aided detection chest radiography: design and results of a randomized, controlled trial. PLoS One 2013; 8(3):e59650.

McWilliams A, et al. Probability of cancer in pulmonary nodules detected on first screening CT. N Engl J Med 2013;69(1):910-9.

National Comprehensive Cancer Network®. (NCCN). Clinical practice guidelines in oncology. Lung cancer screening. Version 2.2019 [https://www.nccn.org/professionals/physician_gls/pdf/lung_screening.pdf] accessed 12/23/18.

Rasmussen JF, et al. Healthcare costs in the Danish randomized controlled lung cancer CT-screening trial: a registry study. Lung Cancer 2014 Mar;83(3):347-55.

Ruano-Ravina A, et al. Lung cancer screening with low-dose computed tomography after the National Lung Screening Trial. The debate is still open. Arch Bronconeumol 2013 Apr;49(4):158-65.

Saghir Z, et al. CT screening for lung cancer brings forward early disease. The randomised Danish Lung Cancer Screening Trial: status after five annual screening rounds with low-dose CT. Thorax 2012;67:296-301.

*Smith RA, et al. American Cancer Society guidelines for the early detection of cancer, 2006. CA Cancer J Clin 2006 Jan-Feb;56(1):11-25.

Medical Policy: LOW-DOSE COMPUTED TOMOGRAPHY (LDCT) FOR LUNG CANCER SCREENING Policy Number: 6.01.19

Page: 6 of 6

*Sobue T, et al. Screening for lung cancer with low-dose helical computed tomography: anti-lung cancer association project. J Clin Oncol 2002 Feb 15;20(4):911-20.

Tanner NT, et al. New testing for lung cancer screening. Oncology (Williston Park). 2012 Feb;26(2):176-82.

U.S. Preventive Services Task Force. Screening for lung cancer. 2013 Dec 31. Update in Progress [https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryDraft/lung-cancer-screening1] accessed 12/23/18.

Wagnetz U, et al. CT screening for lung cancer: implication of lung biopsy recommendations. AJR Am J Roentgenol 2012 Feb;198(2):351-8.

*Key Article

KEY WORDS

EBCT, Electron beam computed tomography, Helical CT, Low-dose CT, Spiral CT.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a Decision Memo for Screening for Lung Cancer with Low Dose Computed Tomography (LDCT). Please refer to the following CMS website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=274>.

There is currently a National Coverage Determination (NCD) for Lung Cancer Screening with Low Dose Computed Tomography (LDCT). Please refer to the following NCD website for Medicare Members: <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=364&ncdver=1&bc=AAAAGAAAAAAAAA%3d%3d&>