

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER
Policy Number	7.01.27
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	10/20/05, 07/20/06, 05/17/07, 05/14/08, 05/28/09, 04/22/10, 06/16/11, 05/24/12, 06/20/13, 05/22/14, 04/16/15, 03/17/16, 03/16/17, 01/18/18, 04/18/19
Archived Date	02/21/02 – 10/20/05
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, Maze procedures, performed on a non-beating heart during cardiopulmonary bypass have been medically proven to be effective and therefore **medically appropriate** for the treatment of medically refractory, chronic, symptomatic atrial fibrillation or flutter, with or without concurrent cardiac surgery.
- II. Based upon our criteria and assessment of the peer-reviewed literature, minimally invasive, off-pump, Maze procedures (e.g., mini thoracotomy), including hybrid or convergent ablation procedures (catheter and surgical ablation), are considered **investigational** as a treatment of atrial fibrillation or flutter

This policy does not address percutaneous transcatheter ablation procedures for the treatment of cardiac arrhythmias.

Refer to Corporate Medical Policy #11.01.03 Experimental and Investigational Services.

POLICY GUIDELINES

The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia, with a prevalence estimated at 0.4% of the population, increasing with age. AF is a supraventricular tachyarrhythmia, characterized by disorganized atrial activation with ineffective atrial ejection. The underlying mechanism of AF involves an interplay between electrical triggering events and the myocardial substrate that permits propagation and maintenance of the aberrant electrical circuit. The most common focal trigger of AF appears to be located within the cardiac muscle that extends into the pulmonary veins. Atrial flutter is considered a variant of AF. Due to the necessity of long-term drug therapy and its associated potential toxicity in patients with AF, surgical techniques have been developed as part of the armamentarium of alternative non-pharmacological treatments. Literature describes patients with drug-resistant AF and flutter as having experienced their arrhythmias for an average of seven years or more and having unsuccessful results with an average of five or more antiarrhythmic medications.

The classic Cox Maze III procedure is a complex surgical procedure that involves sequential atriotomy incisions that interrupt potential re-entrant circuits, which interrupts the aberrant atrial conduction pathways in the heart in cases of AF. It is indicated for patients who do not respond to medical or other surgical antiarrhythmic therapies and is often performed in conjunction with correction of structural conditions of the heart, such as valve repair or replacement. The procedure has

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 2 of 8

become the gold standard technique for the surgical treatment of drug-resistant AF. This procedure is performed on a nonbeating heart during cardiopulmonary bypass.

The Maze procedure entails making incisions in the heart that:

- I. guide an impulse from the sinoatrial (SA) node to the atrioventricular (AV) node;
- II. preserve activation of the entire atrium; and
- III. block re-entrant impulses that are responsible for AF or atrial flutter (AFL).

Despite its high success rate, the traditional “cut and sew” Maze procedure has not been widely utilized other than for those patients who also require concomitant cardiac surgery necessitating the need for cardiopulmonary bypass. Therefore, simplification of the Maze procedure, sometimes referred to as the Cox-Maze IV procedure, has evolved with the use of different ablation tools, such as microwave, cryotherapy, ultrasonography and radiofrequency energy sources to create atrial ablative lesions instead of employing the incisional technique used in the traditional Maze procedure.

Due to the complexity and technical difficulty, associated with the Cox-Maze procedure, less invasive, trans-thoracic, endoscopic, off-pump procedures to treat refractory AF are also being developed and evaluated. Examples of these minimally invasive, off-pump surgical techniques include the thoracoscopic Wolf MiniMaze and the Ex-Maze which uses a paracardioscopy approach.

Studies are also starting to emerge investigating a hybrid approach that combines off-pump surgical and endocardial percutaneous catheter ablation. Hybrid ablation or convergent procedure refers to a procedure that uses both thoracoscopic and percutaneous approaches in the same patient. Ablation is performed on the outer surface of the heart (epicardial) via the thoracoscopic approach, and on the inner surface of the heart (endocardial) via the percutaneous approach. The rationale for doing a hybrid procedure is that a combination of both techniques may result in more complete ablation. Thoracoscopic epicardial ablation is limited by the inability to perform all possible ablation lines, because the posterior portions of the heart are not accessible via thoracoscopy. Percutaneous, endoscopic ablation is limited by incomplete ablation lines that often require repeat procedures. By combining both procedures, a full set of ablation lines can be performed, and incomplete ablation lines can be minimized. This convergent ablation procedure has been proposed for highly symptomatic patients with persistent AF and long-standing persistent AF for whom stand-alone surgical or endocardial ablation procedures have provided unsatisfactory outcomes.

RATIONALE

In January 2002, the FDA approved the Medtronic Cardioblate System, which uses radiofrequency energy to ablate cardiac tissues. The Cardima SAS (surgical ablation system) used during mini-thoracotomy received 510(K) approval by the FDA in 2003 as substantially equivalent to the Medtronic device, amongst others for performing ablation of cardiac tissue during heart surgery via the use of RF energy. Another bipolar RF device used to perform ablations is manufactured by Aticure, Inc. The device has FDA approval for ablation and coagulation of soft tissue during General, ENT, Thoracic, Gynecology & Urology surgical procedures.

Evidence from a number of prospective and retrospective studies conclude that the Maze procedure is effective in restoring sinus rhythm in up to 90% of patients with medically refractory, chronic, symptomatic AF. In addition, there is evidence that, when performed in conjunction with valve repair or replacement, the Maze procedure may reduce the risk of stroke, compared with valve replacement alone (e.g., Reston, et al. 2005, Lim et al. 2010, Budera, et al 2012, Ad, et al. 2013).

There are numerous modifications on the original maze procedure, with variations in the surgical approach, the lesion set used, and the methods for creating lesions (e.g., cut and sew, RFA, etc.). While the evidence on comparative effectiveness of the different approaches is not of high quality, there is evidence from matched case series that indicate that there are not large differences in efficacy among the different approaches (e.g., Khargi, et al. 2005, Stulak, et al. 2007).

While studies have evaluated the minimally invasive, off-pump, epicardial maze procedures, the data are insufficient to reach conclusions about the relative effectiveness of these procedures compared to the classic Maze procedure for the treatment of AF (e.g., Wang, et al. 2011, La Meir, et al. 2013). Some case-series investigating off-pump procedures include only patients who have failed previous catheter ablation. These studies report high success rates following

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 3 of 8

thoracoscopic ablation, suggesting that patients who fail catheter ablation may still benefit from thoracoscopic ablation. However, these series are small and do not provide complete information on comparative efficacy or adverse events (e.g., Okada, et al. 2013).

The Heart Rhythm Society (HRS) (2017) has stated hybrid approach could hold significant promise for those patients with persistent or long-standing persistent, drug-resistant AF to offer improved results over minimal access surgical ablation or catheter ablation alone. It might be reasonable to apply the indications for stand-alone surgical ablation described above to patients being considered for hybrid surgical ablation (Class IIb, LOE C).

There is limited literature related to the use of the hybrid approach in the treatment of AF. While short-term outcomes appear promising, further studies are necessary to determine whether the hybrid approach is effective, especially in patients with long-standing persistent and persistent lone AF (LaMeir, et al. 2012, Pison, et al. 2012, Bisleri, et al. 2013, Gehi, et al. 2013).

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
33254	Operative tissue ablation and reconstruction of atria, limited (e.g., modified maze procedure)
33255 (E/I)	Operative tissue ablation and reconstruction of atria, extensive (e.g., maze procedure); without cardiopulmonary bypass
33256	Operative tissue ablation and reconstruction of atria, extensive (e.g., maze procedure); with cardiopulmonary bypass
33257	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), limited (e.g., modified maze procedure) (List separately in addition to code for primary procedure)
33258 (E/I)	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), extensive (e.g., maze procedure) without cardiopulmonary bypass
33259	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), extensive (e.g., maze procedure) with cardiopulmonary bypass (List separately in addition to code for primary procedure)
33265 (E/I)	Endoscopy, surgical; operative tissue ablation and reconstruction of atria, limited (e.g., modified maze procedure), without cardiopulmonary bypass
33266 (E/I)	Endoscopy, surgical; operative tissue ablation and reconstruction of atria, extensive (e.g., maze procedure), without cardiopulmonary bypass

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No Codes	

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 4 of 8

ICD10 Codes

Code	Description
I48.0-I48.92	Atrial fibrillation and flutter (code range)

REFERENCES

- *Abreu Filho CA, et al. Effectiveness of the maze procedure using cooled-tip radiofrequency ablation in patients with permanent atrial fibrillation and rheumatic mitral valve disease. Circ 2005 Aug 30;112(9 Suppl):I20-5.
- Ad N, et al. Association of operative risk with the outcome of concomitant Cox Maze procedure: a comparison of results across risk groups. J Thorac Cardiovasc Surg 2014 Dec;148(6):3027-33.
- Ad N, et al. Do we increase the operative risk by adding the Cox Maze III procedure to aortic valve replacement and coronary artery bypass surgery? J Thorac Cardiovasc Surg 2012 Apr;143(4):936-44.
- Ad N, et al. Impact of atrial fibrillation duration on the success of first-time concomitant Cox maze procedures. Ann Thorac Surg 2015 Nov;100(5):1613-8.
- Ad N, et al. Minimally invasive stand-alone Cox-maze procedure for patients with nonparoxysmal atrial fibrillation. Ann Thorac Surg 2013 Sep;96(3):792-8.
- Ad N, et al. Minimally invasive stand-alone Cox-maze procedure for persistent and long-standing persistent atrial fibrillation: perioperative safety and 5-year outcomes. Circ Arrhythm Electrophysiol 2017;10:e005352.
- Ad N, et al. The effect of the Cox-maze procedure for atrial fibrillation concomitant to mitral and tricuspid valve surgery. J Thorac Cardiovasc Surg 2013 Dec;146(6):1426-34.
- *Ad N, et al. The outcome of the Cox maze procedure in patients with previous percutaneous catheter ablation to treat atrial fibrillation. Ann Thorac Surg 2011 May;91(5):1371-7.
- Albage A, et al. Early and long-term mortality in 536 patients after the Cox-maze III procedure: a national registry-based study. Ann Thorac Surg 2013 May;95(5):1626-32.
- Badhar V, et al. The Society of Thoracic Surgeons 2017 Clinical Practice Guidelines for the Surgical Treatment of Atrial Fibrillation. Ann Thorac Surg 2017 Jan;103(1):329-341.
- *Barnett SD, AD N. Surgical ablation as treatment for the elimination of atrial fibrillation: a meta-analysis. J Thorac Cardiovasc Surg 2006 May;131(5):1029-35.
- *Ballaux PK, et al. Freedom from atrial arrhythmias after classic maze III surgery: a 10-year experience. J Thorac Cardiovasc Surg 2006 Dec;132(6):1433-40; comment pg 1253.
- *Beukema WP, et al. Intermediate to long-term results of radiofrequency modified maze procedure as an adjunct to open-heart surgery. Ann Thorac Surg 2008 Nov;86(5):1409-14.
- Bisleri G, et al. Hybrid approach for the treatment of long-standing persistent atrial fibrillation: electrophysiological findings and clinical results. Eur J Cardiothorac Surg 2013 Nov;44(5):919-23.
- BlueCross Blue Shield Association. Open and thorascopic approaches to treat atrial fibrillation (Maze and related procedures). Medial Policy Reference Manual Policy #7.01.14. 2018 May 05.
- Boersma LV, et al. Atrial fibrillation catheter ablation versus surgical ablation treatment (FAST): a 2-center randomized clinical trial. Circulation 2012 Jan 3;125(1):23-30.
- Budera P, et al. Comparison of cardiac surgery with left atrial surgical ablation vs. cardiac surgery without atrial ablation in patients with coronary and/or valvular heart disease plus atrial fibrillation: final results of the PRAGUE-12 randomized multicenter study. Eur Heart J 2012 Nov;33(21):2644-52.
- Budera P, et al. Two-staged hybrid ablation of non-paroxysmal atrial fibrillation: clinical outcomes and functional improvements after 1 year. Interact Cardiovasc Thorac Surg 2017 Jul 31 [Epub ahead of print].

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 5 of 8

Calkins H, et al. 2012 HRS/EHRA/ECAS Expert Consensus statement on catheter and surgical ablation of atrial fibrillation: recommendations for patient selection, procedural techniques, patient management and follow-up, definitions, endpoints, and research trial design. Europace 2012 Apr;14(4):528-606.

Calkins H, et al. 2017 HRS/EHRA/ECAS/APHRS/SOLAECE Expert Consensus statement on catheter and surgical ablation of atrial fibrillation. Heart Rhythm. 2017 Oct;14(10):e275-e444.

Chernyavskiy A, et al. Quality of life after surgical ablation of persistent atrial fibrillation: a prospective evaluation. Heart Lung Circ 2016 April;25(4):378-383.

Civello K, et al. Combined endocardial and epicardial ablation for symptomatic atrial fibrillation: single center experience in 100+ consecutive patients. J Innov Cardiac Rhythm Manage 2013;000:1-7.

Cox JL, et al. A hybrid Maze procedure for long-standing persistent atrial fibrillation. Ann Thor Surg 2018 Aug 14 [Epub ahead of print].

Damiano RJ Jr, et al. The CURE-AF trial: a prospective, multicenter trial of irrigated radiofrequency ablation for the treatment of persistent atrial fibrillation during concomitant cardiac surgery. Heart Rhythm 2014 Jan;11(1):39-45.

Davies RA, et al. Surgical and hybrid ablation of atrial fibrillation. Heart Lung Circ 2017 Sep;26(9):960-66.

*Fuster V, et al. ACC/AHA/ESC 2006 Guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association. Circulation 2006 Aug 15;114(7):e257-354.

Gaita F, et al. Very long-term results of surgical and transcatheter ablation of long-standing persistent atrial fibrillation. Ann Thorac Surg 2013 Oct;96(4):1273-8.

*Gaynor SL, et al. A prospective, single-center clinical trial of a modified Cox maze procedure with bipolar radiofrequency ablation. J Thorac Cardiovasc Surg 2004 Oct;128(4):535-42.

*Gaynor SL, et al. Surgical treatment of atrial fibrillation: predictors of late recurrence. J Thorac Cardiovasc Surg 2005 Jan;129(1):104-11.

Gehi AK, et al. Hybrid epicardial-endocardial ablation using a pericardioscopic technique for the treatment of atrial fibrillation. Heart Rhythm 2013 Jan;10(1):22-8.

Gelsomino S, et al. Treatment of lone atrial fibrillation: a look at the past, a view of the present and a glance at the future. Eur J Cardiothorac Surg 2012 Jun;41(6):1284-94.

Genev IK, et al. Comparison of the efficacy and complication rates of the hybrid maze, complete Cox-maze and catheter ablation in the treatment of atrial fibrillation. J Atr Fibrillation 2017 Feb 28;9(5):1543.

Gersak B, et al. Low rate of atrial fibrillation recurrence verified by implantable loop recorder monitoring following a convergent epicardial and endocardial ablation of atrial fibrillation. J Cardiovasc Electrophysiol 2012 Oct;23(10):1059-66.

Gersak B, et al. European experience of the convergent atrial fibrillation procedure: multicenter outcomes in consecutive patients. J Thorac Cardiovasc Surg 2014 Apr;147(4):1411-6.

Geuzebroek GS, et al. Totally thoracoscopic left atrial Maze: standardized, effective and safe. Interact Cardiovasc Thorac Surg 2016 March;22(3):259-264.

Gilligan DM, et al. Multidisciplinary collaboration for the treatment of atrial fibrillation: convergent procedure outcomes from a single center. J Innov Cardiac Rhythm Manage 2013 Oct;1-7.

Gillinov AM, et al. Surgical ablation of atrial fibrillation during mitral-valve surgery. N Engl J Med 2015 Apr 9;372(15):1399-409.

Henn MC, et al. Effectiveness of surgical ablation in patients with atrial fibrillation and aortic valve disease. Ann Thorac Surg 2015 Oct;100(4):1253-9.

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 6 of 8

Huffman MD, et al. Concomitant atrial fibrillation surgery for people undergoing cardiac surgery. *Cochrane Database Syst Rev* 2016 Aug 22;(8):CD011814.

Hwang SK, et al. Long-term outcomes of the maze procedure combined with mitral valve repair: risk of thromboembolism without anticoagulation therapy. *Ann Thorac Surg* 2015 Sep;100(3):840-3.

Ismail I, et al. Left-sided mini-maze procedure via the left atrial appendage. *Interact Cardiovasc Thorac Surg* 2014 Jun;18(6):847-9.

Je HG, et al. A systematic review of minimally invasive surgical treatment for atrial fibrillation: a comparison of the Cox-Maze procedure, beating-heart epicardial ablation, and the hybrid procedure on safety and efficacy. *Eur J Cardiothorac Surg* 2015 Oct;48(4):531-40.

Johansson B, et al. Low mortality and low rate of perceived and documented arrhythmias after Cox maze III surgery for atrial fibrillation. *Pacing Clin Electrophysiol* 2014 Feb;37(2):147-56.

Kurfurst V, et al. Two-staged hybrid treatment of persistent atrial fibrillation: short-term single-centre results. *Interact Cardiovasc Thorac Surg* 2014 Apr;18(4):451-6.

*Khargi K, et al. Surgical treatment of atrial fibrillation; a systematic review. *Eur J Cardiothorac Surg* 2005 Feb;27(2):258-65.

*Kim JB, et al. Long-term outcomes of mechanical valve replacement in patients with atrial fibrillation. Impact of the maze procedure. *Circulation* 2012 May;125(5):2071-80.

La Meir M, et al. Improvement of left atrial function and left atrial reverse modeling after minimally invasive radiofrequency ablation evaluated by two-dimensional speckle tracking echocardiography. *J Thorac Cardiovasc Surg* 2013;146(1):72-77.

La Meir M, et al. Minimal invasive surgery for atrial fibrillation: an updated review. *Europace* 2013 Feb;15(2):170-82.

La Meir M, et al. Minimally invasive surgical treatment of lone atrial fibrillation: early results of hybrid versus standard minimally invasive approach employing radiofrequency sources. *Int J Cardiol* 2013 Aug 20;167(4):1469-75.

Lawrance CP, et al. Comparison of the stand-alone Cox-Maze IV procedure to the concomitant Cox-Maze IV and mitral valve procedure for atrial fibrillation. *Ann Cardiothorac Surg* 2014 Jan;3(1):55-61.

Lawrance CP, et al. A minimally invasive Cox maze IV procedure is as effective as sternotomy while decreasing major morbidity and hospital stay. *J Thorac Cardiovasc Surg* 2014 Sep;148(3):955-61.

Lawrance CP, et al. Surgical ablation for atrial fibrillation: techniques, indications, and results. *Curr Opin Cardiol* 2015 Jan;30(1):58-64.

Lawrance CP, et al. Surgery for atrial fibrillation. *Heart Fail Clin* 2016 April;12(2):235-243.

Massimiano PS, et al. Minimally invasive fibrillating heart surgery: a safe and effective approach for mitral valve and surgical ablation for atrial fibrillation. *Ann Thorac Surg* 2013 Aug;96(2):520-7.

McKinnie J. The convergent procedure- a standardized and anatomic approach addresses the clinical and economic unmet needs of the persistent atrial fibrillation population. *Arrhythmia Electrophysiol Rev* 2013;2(2):145-8.

Mei J, et al. Complete thoracoscopic ablation of the left atrium via the left chest for treatment of lone atrial fibrillation. *J Thorac Cardiovasc Surg* 2014 Jan;147(1):242-246.

Muneretto C, et al. Durable staged hybrid ablation with thoracoscopic and percutaneous approach for treatment of long-standing atrial fibrillation: a 30 month assessment with continuous monitoring. *J Thorac Cardiovasc Surg* 2012 Dec;144(6):1460-5.

Muneretto C, et al. Successful treatment of lone persistent atrial fibrillation by means of a hybrid thoracoscopic-transcatheter approach. *Innovations* 2012 Jul-Aug;7(4):254-8.

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 7 of 8

- Musharbash FN, et al. Performance of the Cox-maze IV procedure is associated with improved long-term survival in patients with atrial fibrillation undergoing cardiac surgery. J Thorac Cardiovasc Surg 2018Jan;155(1):159-170.
- *Nakajima H, et al. The effect of cryo-maze procedure on early and intermediate term outcome in mitral valve disease: case matched study. Circ 2002 Sep 24;106(12 Suppl 1):I46-50.
- Okada S, et al. The impact of previous catheter-based ablation on the efficacy of the Cox-maze IV procedure. Ann Thorac Surg 2013 Sep;96(3):786-91.
- Phan K, et al. Effectiveness and safety of simultaneous hybrid thoracoscopic endocardial catheter ablation of atrial fibrillation in obese and non-obese patients. J Thorac Dis 2017;9(9):3087-3096.
- Phan K, et al. Systematic review and meta-analysis of surgical ablation for atrial fibrillation during mitral valve surgery. Ann Cardiothorac Surg 2014 Jan;3(1):3-14.
- Phan K, et al. Surgical ablation for treatment of atrial fibrillation in cardiac surgery: a cumulative meta-analysis of randomized controlled trials. Heart 2014 May;100(9):722-730.
- Phan K, et al. Thoracoscopic surgical ablation versus catheter ablation for atrial fibrillation. Eur J Cardiothorac Surg 2016 April;49(4):1044-1051.
- Pison L, et al. Hybrid thoracoscopic surgical and transvenous catheter ablation of atrial fibrillation. J Am Coll Cardiol 2012 Jul 3;60(1):54-61.
- Pison L, et al. Effectiveness and safety of simultaneous hybrid thoracoscopic and endocardial catheter ablation of lone atrial fibrillation. Ann Cardiothorac Surg 2014 Jan;3(1):38-44.
- Pokushalov E, et al. catheter versus surgical ablation of atrial fibrillation after a failed initial pulmonary vein isolation procedure: a randomized controlled trial. J Cardiovasc Electrophysiol 2013 Dec;24(12):1338-43.
- Richardson TD, et al. Staged versus simultaneous thoracoscopic hybrid ablation for persistent atrial fibrillation does not affect time to recurrence of atrial arrhythmia. J Cardiovasc Electrophysiol 2016 April;27(4):428-434.
- Ruaengsri C, et al. The Cox-Maze IV procedure in its second decade: still the gold standard? Eur J Cardiothorac Surg 2018 Apr 1;53(suppl_1):i19-i25.
- Saint LL, et al. Cox-Maze IV results for patients with lone atrial fibrillation versus concomitant mitral disease. Ann Thorac Surg 2012 Mar;93(3):789-94.
- Saint LL, et al. Incremental risk of the Cox-maze IV procedure for patients with atrial fibrillation undergoing valve surgery. J Thorac Cardiovasc Surg 2013 Nov;146(5):1072-7.
- Sirak JH, et al. Interim results of the 5-box thoracoscopic maze procedure. Ann Thorac Surg 2012 Dec;94(6):1880-4.
- Stulak JM, et al. Surgical ablation for atrial fibrillation for two decades: Are results of the new techniques equivalent to the Cox maze III procedure? J Thorac Cardiovasc Surg 2014 May;147(5):1478-1486.
- Thosani AJ, et al. Closed chest convergent epicardial-endocardial ablation of non-paroxysmal atrial fibrillation- a case series and literature review. Arrhythmia Electrophysiol Rev 2013;2(1):65-8.
- Van Laar C, et al. The totally thoracoscopic maze procedure for the treatment of atrial fibrillation. Interact Cardiovasc Thorac Surg 2017 Jan;24(1):102-111.
- Velagic V, et al. Repeat procedures after hybrid thoracoscopic ablation in the setting of longstanding persistent atrial fibrillation: electrophysiological findings and 2-year clinical outcome. J Cardiovasc Electrophysiol 2016 Jan;27(1):41-50.
- Watkins AC, et al. Prospective assessment of the cryomaze procedure with continuous outpatient telemetry in 136 patients. Ann Thorac Surg 2014 Apr;97(4):1191-8.
- Weimar T, et al. The cox-maze procedure for lone atrial fibrillation: a single center experience over 2 decades. Circ Arrhythm Electrophysiol 2012 Feb;5(1):8-14.

Medical Policy: MAZE PROCEDURES FOR ATRIAL FIBRILLATION AND FLUTTER

Policy Number: 7.01.27

Page: 8 of 8

Weimar T, et al. Approaching a paradigm shift: endoscopic ablation of lone atrial fibrillation on the beating heart. Ann Thorac Surg 2012 Dec;94(6):1886-92.

Yanagawa B, et al. Outcome of concomitant Cox-maze III procedure using argon-based cryosurgical system: a single-center experience with 250 patients. Ann Thorac Surg 2013 May;95(5):1633-9.

Zembala M, et al. Minimally invasive hybrid ablation procedure for the treatment of persistent atrial fibrillation: one year results. Kardiologia Polska 2012;70(8):819-28.

*Key Article

KEY WORDS

AF, Atrial fibrillation, Atrial Flutter, MAZE, Convergent procedure, COX-III, Epicardial Maze, Ex-Maze, Hybrid, MiniMaze, Thoracoscopic off-pump surgical ablation (TOPS).

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based on our review, the Maze procedure as treatment for atrial fibrillation is not addressed in National or Regional Medicare coverage determinations or policies. “Though no formal coverage determination exists, Medicare may provide coverage without a formal coverage determination.”