

MEDICAL POLICY

| MEDICAL POLICY DETAILS | |
|------------------------|---|
| Medical Policy Title | POSTUROGRAPHY |
| Policy Number | 2.01.20 |
| Category | Technology Assessment |
| Effective Date | 10/18/01 |
| Revised Date | 05/16/02, 04/24/03 |
| Archived Date | 02/19/04 |
| Edited Date | 11/10/05, 12/21/06, 12/20/07, 12/18/08, 12/17/09, 12/16/10, 12/15/11, 12/20/12, 12/19/13, 12/18/14, 12/17/15, 12/15/16, 12/21/17, 12/20/18 |
| Product Disclaimer | <ul style="list-style-type: none"> • <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i> • <i>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</i> • <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i> |

POLICY STATEMENT

Based upon our criteria and assessment of the peer-reviewed literature, posturography does not improve patient outcomes and therefore, is considered to be not medically necessary.

POLICY GUIDELINES

The Federal Employees Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus, these procedures, devices, or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Posturography is a diagnostic test used to assess balance problems that may be caused by peripheral vestibular or oculovestibular dysfunction. Also known as moving platform posturography, dynamic posturography or Equitest™, the test involves the use of a platform that can be manipulated to assess the patient's responses to various demands on postural equilibrium control (balance). It attempts to identify disequilibrium due to pathology of the labyrinthine sensory organs (e.g., semicircular canals and otoliths) and the oculovestibular and somatosensory vestibular pathways. It is proposed that posturography is useful to assess progress in patients undergoing rehabilitation for balance disorders and postural deficits.

RATIONALE

Girardi and colleagues evaluated the results of computerized dynamic posturography in a group of 65 elderly patients who had a history of falling. While 78% of patients had abnormal results on computerized dynamic posturography testing, results in a broader group of patients is required to determine whether or not results of computer dynamic posturography can identify those at risk for falling. Baloh and colleagues studied balance control in a group of elderly patients who complained of balance disorders and a group of age-matched controls; the subjects were tested in a variety of situations (e.g., eyes open and closed, while standing on a foam pad to disrupt sensory cues, tilting of the platform, etc.). The authors concluded that posturography data provided little information about the cause of imbalance and did not correlate with the frequency of reported falls. Other authors have pointed out that how results of computerized dynamic posturography may correlate to functional activities, such as gait, is uncertain. For example, measurements of gait, frequently gait velocity, are often used in the elderly to assess balance and mobility. A variety of patient questionnaires have been designed to measure self-perceived dizziness or balance. The correlation between the results of these clinical tests, questionnaires, and computerized dynamic posturography are uncertain.

Medical Policy: POSTUROGRAPHY

Policy Number: 2.01.20

Page: 2 of 3

The contribution of dynamic posturography to the overall assessment and customization of the exercise program is unclear. In particular, the reports do not describe how (or whether) the exercise programs were modified based on specific deficits identified by platform posturography. There are no controlled trials have assessed whether individually customized therapy programs are more effective than generic vestibular exercises. The literature has not demonstrated that posturography provides any improvement in health outcomes over standard physical therapy; it has not been shown to be clinically appropriate for diagnosis or treatment; nor has it demonstrated improved treatment decision making and/or health outcomes based on the results of dynamic posturography.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- *CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.*
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*
- *Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).*

CPT Codes

| Code | Description |
|-------------|------------------------------------|
| 92548 (NMN) | Computerized dynamic posturography |

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

| Code | Description |
|-------------|--------------------|
| | No specific code |

ICD10 Codes

| Code | Description |
|-----------------|---|
| H81.10-H81.13 | Benign paroxysmal vertigo (code range) |
| H81.20-H81.23 | Vestibular neuronitis (code range) |
| H81.311-H81.319 | Aural vertigo (code range) |
| H81.391-H81.399 | Other peripheral vertigo (code range) |
| H81.8x1-H81.93 | Other disorders of vestibular function (code range) |
| H82.1-H82.9 | Vertiginous syndromes in diseases classified elsewhere (code range) |
| H83.2x1-H83.2x9 | Labyrinthine dysfunction (code range) |
| H83.8x1-H83.8x9 | Other specified diseases of inner ear (code range) |
| H83.90-H83.93 | Unspecified disease of inner ear (code range) |

REFERENCES

Amin M, et al. A comparison of electronystagmography results with posturography findings from the balance Trak 500. Otol Neurotol 2002 Jul;23(4):488-93.

Baloh RW, et al. Balance disorders in older persons: quantification with posturography. Otolaryngol Head Neck Surg 1998; 119(1):89-92.

Black FO. What can posturography tell us about vestibular function? Ann NY Acad Sci 2001 Oct;942:446-64.

BlueCross BlueShield Association. Dynamic posturography. Medical Policy Reference Manual Policy #2.01.02. 2017 Feb 9.

Medical Policy: POSTUROGRAPHY

Policy Number: 2.01.20

Page: 3 of 3

Clendaniel RA. Outcome measures for assessment of treatment of the dizzy and balance disorder patient. Otolaryngol Clin North Am 2000; 33(3):519-33.

Evans MK, et al. Posturography does not test vestibulospinal function. Otolaryngol Head Neck Surg 1999 Feb;120(2):164-73.

Gianoli G, et al. Posturographic performance in patients with the potential for secondary gain. Otolaryngol Head Neck Surg 2000 Jan;122(1):11-8.

Girardi M, et al. Predicting fall risks in an elderly population: computer dynamic posturography versus electronystagmography test results. Laryngoscope 2001 Sep;111(9):1528-32.

Harstall C. Computerized dynamic posturography in rehabilitation: has its reliability and validity been established? Physiother Can 2000 Winter;56-63.

Ondo W, et al. Computerized posturography analysis of progressive supranuclear palsy. Arch Neurol 2000 Oct;57:1464-9.

Reid VA, et al. Using posturography to detect unsteadiness in 13 patients with peripheral neuropathy: a pilot study. Neurol Clin Neurophysiol 2002;4:2-8.

Rose DJ, et al. Can the control of bodily orientation be significantly improved in a group of older adults with a history of falls?. J Am Geriatr Soc 2000 Mar;48(3):275-82.

Ruckenstein MJ, et al. Balance function testing: a rational approach. Otolaryngol Clin N Amer 2000 Jun;33(3):507-7.

*Key Article

KEY WORDS

Balance, Equitest, Posturography, Vestibular dysfunction.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based on our review, posturography is not addressed in National or Regional Medicare coverage determinations or policies.