

MEDICAL POLICY

| MEDICAL POLICY DETAILS | |
|------------------------|--|
| Medical Policy Title | VISION THERAPY |
| Policy Number | 9.01.04 |
| Category | Therapy/Rehabilitation |
| Effective Date | 10/18/01 |
| Revised Date | 04/24/03, 02/19/04, 02/17/05, 11/17/05, 02/22/07, 12/13/07 |
| Archived Date | 12/11/08 |
| Edited Date | 12/10/09, 12/09/10, 12/08/11, 12/06/12, 10/24/13, 12/11/14, 12/10/15, 12/8/16, 12/14/17, 12/13/18 |
| Product Disclaimer | <ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit. |

POLICY STATEMENT

- I. Based upon our criteria and assessment of peer-reviewed literature, vision therapy that includes orthoptics and occlusion therapy, has been medically proven to be effective and is considered **medically appropriate** for the treatment of *amblyopia*.
- II. Based upon our criteria and assessment of peer-reviewed literature, vision therapy that includes prism adaptation, has been medically proven to be effective and is considered **medically appropriate** when utilized for acquired *esotropia* prior to surgical intervention.
- III. Based upon our criteria and assessment of peer-reviewed literature, vision therapy has been medically proven to be effective and is considered **medically appropriate** for the treatment of *convergence insufficiency*.
- IV. Based upon our criteria and the lack of peer-reviewed literature, vision therapy has not been medically proven to be effective and is considered **investigational**, including but not limited to, the following indications:
 - A. All other accommodative and vergence dysfunctions such as: fusional vergence dysfunction, divergence excess, convergence excess, divergence insufficiency, vertical phorias, basic exophoria, basic esophoria, accommodative insufficiency, sustained accommodation, accommodative infacility and spasm accommodation (see statement III above on coverage of convergence insufficiency).
 - B. Low vision;
 - C. Myopia;
 - D. Nystagmus;
 - E. Presbyopia;
 - F. Strabismus, including esotropia (with the exception of acquired esotropia as stated above), and exotropia; and
 - G. Age-related macular degeneration.
- V. Based upon our criteria and assessment of the peer-reviewed literature, vision therapy does not improve patient outcomes and is **investigational** for *learning disabilities*; including Attention Deficit Hyperactivity Disorder (ADHD) and dyslexia.

DESCRIPTION

Vision therapy (also known as visual therapy, visual training, vision training, eye training) involves a range of treatment modalities that includes the use of lenses, prisms, filters, optometric phototherapy (Syntonics), occlusion therapy (eye patching), behavioral modalities, and eye exercises (orthoptics, pleoptics). The therapeutic goal of vision therapy is to

Medical Policy: VISION THERAPY

Policy Number: 9.01.04

Page: 2 of 4

correct or improve specific visual dysfunctions. Dysfunctions that purportedly are treatable by vision therapy include amblyopia, strabismus, binocular, accommodative and convergence disorders as well as reading disorders such as dyslexia that are thought to be related to a lack of eye coordination. Vision therapy is performed in an optometrist’s or ophthalmologist’s office 1-2 times weekly for a number of months with additional home exercises done as reinforcement.

RATIONALE

Most studies evaluating the efficacy of vision therapy for visual disorders are small. In general, these studies are poorly designed with significant methodological flaws, and the data derived from them are relatively weak and inconclusive. There is some evidence to support the use of vision therapy that involves occlusion as a treatment for amblyopia (treatment success with patching 72.3% -79%) and vision therapy that involves prism adaptation prior to surgery administered as a treatment for acquired esotropia (surgical success rates for prism adaption prior to surgery 89% versus 72% for patients receiving no prism therapy). Large, well-designed studies comparing vision therapy with other treatment modalities, standardization of outcome measurements and the criteria for defining patient selection criteria are needed to evaluate vision therapy for visual dysfunctions adequately.

A number of optometrists advocate vision therapy for patients with learning disabilities including dyslexia claiming that while vision therapy does not treat these disorders directly, it may improve visual efficiency and visual processing to allow the individual to be more responsive to educational instruction. This rationale for the use of vision therapy as a treatment for reading disabilities is unproven. While research suggests a relationship between oculomotor efficiency and reading skills, other studies have found that reading skills are related to language skills, and oculomotor ability is not the principal cause of reading disability. There is a scarcity of quality data on the efficacy of vision therapy for treating dyslexias and other reading and learning disabilities. Most of these study results were found to be inconsistent and the studies themselves were flawed by serious design limitations (e.g. small sample sizes, poorly defined patient criteria).

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.*
- *CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.*
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

| Code | Description |
|-------------|--|
| 92065 | Orthoptic and/or pleoptic training, with continuing medical direction and evaluation |

Copyright © 2019 American Medical Association, Chicago, IL

ICD10 Codes

| Code | Description |
|-----------------|--|
| H51.11-H52.12 | Convergence insufficiency and excess (code range) |
| H53.001-H53.039 | Amplyopia (code range) |
| | Remainder of diagnosis codes considered investigational or not medically necessary |

REFERENCES

Al-Karmi R, et al. Image relocation with prisms in patients with age-related macular degeneration. Can J Ophthalmol 2006 Jun;41(3):313-8.

American Academy of Ophthalmology. Preferred practice Pattern. Care of the patient with strabismus: esotropia and exotropia. [http://one.aao.org/guidelines-browse?filter=preferredpracticepatterns] accessed 10/20/17.

Medical Policy: VISION THERAPY

Policy Number: 9.01.04

Page: 3 of 4

American Academy of Ophthalmology. Preferred Practice Pattern. Care of the patient with amblyopia. [<http://one.aao.org/guidelines-browse?filter=preferredpracticepatterns>] accessed 10/20/17.

American Academy of Ophthalmology. Policy statement: learning disabilities, dyslexia and vision therapy. [<http://one.aao.org/CE/PracticeGuidelines/ClinicalStatements.aspx?sid=d9939a8b-1675-4bf8-85fb-154652305795>] accessed 10/20/17.

American Academy of Ophthalmology. Joint Technical Report: Learning Disabilities, Dyslexia, and Vision A Joint Technical Report of the AAP, AAO, AAPOS, and AACO on learning disabilities, dyslexia, and vision. [<http://www.aao.org/about/policy/upload/Learning-Disabilities-Dyslexia-Vision-2009.pdf>] accessed 10/20/17.

American Academy of Optometry and the American Optometric Association. Vision therapy. Information for health care and other allied professionals. A joint organizational policy statement of the American Academy of Optometry and the American Optometric Association. [<http://www.aoa.org/documents/JOPS%20on%20Vision%20Therapy.pdf>] accessed 10/20/17.

*American Academy of Pediatrics. Clinical practice guideline: treatment of the school-aged child with attention deficit/hyperactivity disorder. Clinical practice guideline. 2001 Oct;108(4):1033-44.

American Optometric Association. Clinical practice guideline. Care of the patient with accommodative and vergence dysfunction. [<http://www.aoa.org/documents/optometrists/CPG-18.pdf>]. reviewed 2010. accessed 10/20/17.

American Optometric Association. Clinical practice guideline. Care of the patient with amblyopia. [<http://www.aoa.org/documents/optometrists/CPG-4.pdf>] reviewed 2004. accessed 10/20/17.

American Optometric Association. Clinical practice guideline. Care of the patient with learning related vision problems. [<http://www.aoa.org/documents/optometrists/CPG-20.pdf>] reviewed 2006. accessed 10/20/17.

American Optometric Association. Clinical practice guideline. Care of the patient with strabismus. [<http://www.aoa.org/documents/optometrists/QRG-12.pdf>] reviewed 2010. accessed 10/20/17.

Aziz S, et al. Are orthoptic exercises an effective treatment for convergence and fusion deficiencies? Strabismus 2006 Dec;14(4):183-9.

BlueCross BlueShield Association. Orthoptics for the treatment of learning disabilities. Medical Policy Reference Manual Policy # 9.03.03. 2017 March 9.

*Bouldoukian J, et al. Randomized controlled trial of the effect of colored overlays on the rate of reading of people with specific learning disabilities. Ophthalmic Physiol Opt 2002 Jan;22(1):55-60.

Bowers AR, et al. Community-based trial of a peripheral prism visual field expansion device for hemianopia. Arch Ophthalmol 2008 May;126(5):657-64.

Brautaset RL, et al. Effects of orthoptic treatment on the CA/C and AC/A ratios in convergence insufficiency. Invest Ophthalmol Vis Sci 2006 Jul;47(7):2876-80.

Brautaset R, et al. Accommodation insufficiency in children: are exercises better than reading glasses? Strabismus 2008 Apr-Jun;16(2):65-9.

El-Ghrably IA, et al. Does compliance with amblyopia management improve following supervised occlusion treatment? Eur J Ophthalmol 2007 Sep-Oct;17(5):823-7.

Figueira EC, et al. Intermittent exotropia: comparison of treatments. Clin Experiment Ophthalmol 2006 Apr;34(3):245-51.

*Gallaway M, et al. The effectiveness of pencil pushups treatment for convergence insufficiency: a pilot study. Optom Vis Sci 2002 Apr;79(4):265-67.

Helveston EM. Visual training: current status in ophthalmology. Am J Ophthalmol 2005 Nov;140(5):903-10.

*Leiba H, et al. Long-term follow-up of occlusion therapy in amblyopia. Ophthalmol 2001 Sep;108(9):1552-5.

Medical Policy: VISION THERAPY

Policy Number: 9.01.04

Page: 4 of 4

Loudon SE, et al. predictors and a remedy for noncompliance with amblyopia therapy in children measured with the occlusion dose monitor. Invest Ophthalmol Vis Sci 2006 Oct;47(10):4393-400.

Markowitz SN. Principles of modern low vision rehabilitation. Can J Ophthalmol 2006 Jun;41(3):289-312.

Pediatric Eye Disease Investigator Group, et al. A randomized trial of atropine vs patching for treatment of moderate amblyopia: follow-up at age 10 years. Arch Ophthalmol 2008 Aug;126(8):1039-44.

Rubab S, et al. Glued patches for children resistant to amblyopia occlusion therapy. Arch Ophthalmol 2008 Jan;126(1):133-4.

*Scheiman M, et al. A survey of treatment modalities for convergence insufficiency. Optom Vis Sci 2002 Mar;79(3):151-7.

Scheiman M, et al. A randomized clinical trial of treatments for convergence insufficiency in children. Arch Ophthalmol 2005 Jan;123(1):14-24.

Scheiman M, et al. Randomized trial of treatment of amblyopia in children aged 7 to 17 years. Arch Ophthalmol 2005 Apr;123(4):437-47.

Scheiman M, et al. A randomized clinical trial of vision therapy/ orthoptics versus pencil pushups for the treatment of convergence insufficiency in young adults. Optom Vis Sci 2005 Jul;82(7):583-95.

Schotton K, et al. Interventions of strabismic amblyopia. Cochrane Database Syst Rev 2008 Apr 16;(2):CD006461.

Smith HJ, et al. A randomized controlled trial to determine the effectiveness of prism spectacles for patients with age-related macular degeneration. Arch Ophthalmol 2005 Aug;123(8):1042-50.

The National Eye Institute. Amblyopia clinical studies database. [<http://www.nei.nih.gov/health/amblyopia/>] accessed 10/20/17.

*The Pediatric Eye Disease Investigator Group. A randomized trial of atropine vs patching for treatment of moderate amblyopia in children. Arch Ophthalmol 2002 Mar;120:268-78.

*Key Article

KEY WORDS

Acquired esotropia, Amblyopia, Convergence insufficiency, Orthoptics.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based on our review, there is no specific regional or national coverage determination for vision therapy.