

# MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON
Policy Number	1.01.53
Category	Equipment/Supplies
Effective Date	10/18/01
Revised Date	02/21/02, 01/16/03, 02/19/04, 02/24/05, 02/23/06, 12/07/06, 10/24/07, 08/28/08, 10/28/09, 04/28/11, 04/26/12, 06/28/12, 04/25/13, 04/24/14, 04/23/15, 06/22/16, 06/22/17, 04/26/18, 02/28/19
Product Disclaimer	<ul style="list-style-type: none"> <li>• If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</li> <li>• If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</li> <li>• If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</li> </ul>

## POLICY STATEMENT

### I. ELECTRICAL BONE GROWTH STIMULATOR

- A. Based upon our criteria and assessment of the peer-reviewed literature, electrical bone growth stimulation has not been demonstrated to improve patient outcomes in the applications of fresh fractures, stress fractures, *delayed unions*, or fresh bunionectomies and, therefore, is considered **investigational**.
- B. Based upon our criteria and review of the peer reviewed literature *non-invasive* electrical bone growth stimulation for the treatment of *non-union* secondary to trauma has been medically proven to be effective and therefore **medically appropriate** for the following indications:
1. The treatment for *non-union* secondary to trauma of the bones of the appendicular skeleton, including the humerus, ulna, radius, carpals, metacarpals, femur, tibia, fibula, tarsals, metatarsals, phalanges, scapula, clavicle, pelvis and patella. In order for coverage to be available, patients must meet all of the following criteria:
    - a. Greater than or equal to 3 months have elapsed since injury or initial treatment;
    - b. Serial radiographs of the preceding 3 month period have confirmed that no progressive signs of healing have occurred or unless the injury is greater than 6 months, shows no progressive signs of healing, and has not been actively treated;
    - c. The fracture gap is one centimeter or less; and
    - d. The patient can be adequately immobilized and, when appropriate, is likely to comply with non-weight bearing.
  2. The treatment of infantile non-union;
  3. The treatment of failed joint fusion secondary to failed arthrodesis of the ankle or knee; and
  4. As a non-surgical salvage for pseudoarthrosis (minimum nine months after last lumbar spinal fusion surgery).
- C. Contraindications to the use of an electrical bone growth stimulation include:
1. Fracture gaps greater than one centimeter; and
  2. Patients with a demand-type pacemaker or an implantable cardioverter defibrillator.

### II. ULTRASONIC BONE GROWTH STIMULATOR –

- A. Based upon our criteria and assessment of the peer reviewed literature, ultrasound accelerated fracture healing systems have been proven to be medically effective when used to treat *non-union* fractures (excluding fractures of the skull or vertebrae and tumor-related fractures) and are therefore **medically appropriate** when all of the following criteria are met:
1. At least 3 months have elapsed since injury;

## Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON

Policy Number: 1.01.53

Page: 2 of 7

2. Nonunion of the fracture is documented by a minimum of two sets of radiographs obtained prior to starting treatment with the US device, separated by a minimum of 90 days, each including multiple views of the fracture site, and with written interpretation by a physician stating that there has been no clinically significant evidence of fracture healing between the two sets of radiographs;
  3. The fracture gap is one cm or less; and
  4. The patient can be adequately immobilized and is of an age where likely to comply with non-weightbearing.
- B. Based upon our criteria and review of the peer-reviewed literature ultrasound accelerated fracture healing systems do not significantly improve patient outcomes and are therefore **not medically necessary** for the following indications:
1. To accelerate healing of fresh, closed, posteriorly displaced distal radius fractures,
  2. To accelerate healing of fresh, closed or Grade 1 tibial diaphysis fractures;
  3. To accelerate fresh fractures, fusions, or delayed unions of the scaphoid (carpal navicular).
  4. To treat delayed union of fractures;
  5. Treatment of congenital pseudoarthrosis;
  6. Treatment of Charcot arthropathy; (except that treatment of fractures related to Charcot arthropathy using ultrasonic bone growth stimulators are considered medically necessary when all of the criteria listed in IIA are met; and
  7. Treatment of Osteogenesis Imperfecta.

*Refer to Corporate Medical Policy #7.01.40 regarding Bone Growth Stimulators; Invasive and Noninvasive Electrical Stimulation of the Spine.*

*Refer to Corporate Medical Policy # 2.01.31 regarding Extracorporeal Shock Wave Therapy (ESWT) for Musculoskeletal Conditions and Soft Tissue Wounds..*

### **POLICY GUIDELINES**

- I. Prior authorization is contract dependent. Please refer to your Customer (Member/Provider) Services Department for contract information.
- II. Durable Medical Equipment rider/coverage is required.
- III. Ultrasound accelerated healing devices are not to be used in conjunction with any other noninvasive osteogenic stimulation devices.
- IV. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

### **DESCRIPTION**

- I. **Electrical bone growth stimulators** are used to induce the growth of bones in cases of delayed union or non-union of fractures. Two methods of electrical bone growth stimulation are available:
  - A. **Non-invasive** stimulators use an external power supply and externally applied coils that produce an electrical current to the fracture site via pulsed electromagnetic fields (PEMFs), combined electromagnetic field (CEMF) technology, or capacitive coupling to stimulate bone growth.
  - B. **Invasive stimulators** use a current generator that is surgically implanted in an intramuscular subcutaneous space and connected to an electrode that is implanted within the bone fragments that are hoped to be fused. The power source is removed in a second surgical procedure once it has discharged.
- II. **Ultrasonic Accelerated Fracture**, or **Sonic Accelerated Fracture Healing System (SAFHS)**, is a non-invasive device that uses low intensity, pulsed, ultrasound therapy to stimulate and accelerate fracture healing time. The device consists of two main components: a signal generator about the size of a laptop computer and a small, square transducer connected to the generator by cable. The transducer is applied to the skin over the fracture site using a gel to facilitate transmission of the ultrasound signal.

## **Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON**

**Policy Number: 1.01.53**

**Page: 3 of 7**

*Delayed unions* are defined by using clinical and radiographic findings suggesting an un-united fracture where the possibility of healing exists. Healing has not advanced at the "average" rate for the location and type of fracture.

*Non-unions* are defined as radiographic findings with clinical mobility of the bone fragments and where bone healing has ceased and there has been more than 3 months since the time of the fracture.

Delayed union differs from non-union in that in the former, there are no indications that union will fail, while in the latter, there are no longer any visible signs that union will occur.

### **RATIONALE**

The FDA has given premarket approval for the EBI Bone Healing System, the Orthologic Bone Growth Stimulator, SpinalPak, Spinal-Stim Lite, Physio-Stim Life, OrthoPak, and SpinaLogic external stimulators and the Orthogen/Osteogen, Zimmer Direct Current Bone Growth Stimulator, and SpF implanted spinal fusion stimulators.

There is sufficient evidence reported in the peer-reviewed literature to conclude that external electrical stimulation improves outcomes for non-union of fractures, for infantile non-union, failed joint fusion, and for non-surgical salvage for pseudoarthrosis. Non-invasive and invasive electrical bone stimulation improves outcomes when used as an adjunct to spinal fusion surgery for patients at high risk for pseudoarthrosis. Improved outcomes have been achieved outside the investigational setting. A randomized controlled trial to determine if interferential current could significantly reduce healing time in new fractures of the tibia or prevent non-union found no significant difference in time to union compared to placebo. A randomized controlled trial to determine if interferential current would accelerate tibial stress fracture healing found no difference in time to healing between treatment and placebo groups. Greater device use and less weightbearing loading enhanced the effectiveness of the active device. A 2002 meta-analysis of trials of the effect of electrical stimulation on musculoskeletal systems included four studies of fresh fractures, all of them failing to provide evidence of efficacy.

The FDA approved the BioniCare® Stimulator Model BIO-1000™ in 2003 for use as an adjunctive therapy in reducing the level of pain and symptoms associated with osteoarthritis of the knee. The BioniCare® device is purported to stimulate chondrogenesis, however no studies have been performed in humans to evaluate whether chondrogenesis occurs with use of this device. No studies of the use of electrical bone growth stimulators in bunionectomies were identified.

FDA premarket approval was granted the Exogen 2000 Sonic Accelerated Fracture Healing System (SAFHS®) in 1994 for treatment of fresh Colles fractures and open tibial diaphysis fractures when managed by closed reduction and casting and expanded to non-unions in 2000. Data presented to the FDA as part of the approval process for the SAFHS® device demonstrated that 64 of 74 cases of non-union (mean fracture age nearly 3 years) were healed with use of low-intensity ultrasound. Patients receiving drugs which alter bone metabolism were excluded from studies of the device. Two studies of ultrasound after intramedullary nailing and fixation with absorbable screws showed no benefit from ultrasound. Most fresh fractures heal following standard care, such as closed reduction and casting. The United Kingdom's National Institute for Health and Clinical Excellence (NICE) updated their guidance on low-intensity pulsed ultrasound for the treatment of nonunion and delayed fracture healing in 2013. NICE reached the following conclusions: Clinical evidence shows a high rate of fracture healing which supports the use of the EXOGEN ultrasound bone healing system to treat long-bone fractures with nonunion (failure to heal after 9 months). In addition, the EXOGEN ultrasound bone healing system to treat long-bone fractures with nonunion is associated with an estimated cost saving when compared with current management, through avoiding surgery. There is some radiological evidence of improved healing when the EXOGEN ultrasound bone healing system is used for long-bone fractures with delayed healing (no radiological evidence of healing after approximately 3 months). There are substantial uncertainties about the rate at which bone healing progresses without adjunctive treatment between 3 and 9 months after fracture, and about whether or not surgery would be necessary. These uncertainties result in a range of cost consequences, some cost-saving and others that are more costly than current management.

Small randomized controlled trials suggest that ultrasound may accelerate healing of fresh fractures and promote healing in subgroups at risk for non-union. Larger trials are needed to confirm this. No studies were identified that included children less than 17 years old. The mechanism for the effect of ultrasound on bone healing is not fully understood.

## Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON

Policy Number: 1.01.53

Page: 4 of 7

Foley et al published results of the investigational device exemption study of pulsed electromagnetic field (PEMF) stimulation (Cervical-Stim device from Orthofix) as an adjunct to anterior cervical discectomy and fusion (ACDF) with anterior cervical plates and allograft interbody implants. A total of 323 patients were randomized, 163 to PEMF and 160 to no stimulation. All patients were active smokers (more than one pack of cigarettes per day, 159 patients) or were undergoing multilevel ACDF (192 patients). The patients in the treatment group wore the Cervical-Stim device for 4 hours per day for 3 months starting 1 week after surgery. Efficacy was measured by radiographic analysis at 1, 2, 3, 6, and 12 months. Fusion rates for the 240 evaluable patients at 6 months were 83.6% for the PEMF group and 68.6% for the control group (p=.0065). By intent-to-treat analysis, assuming that nonevaluable patients did not have fusion, PEMF and control groups fusion rates were 65.6% and 56.3%, respectively (p=.0835). Of 245 patients available for follow-up at 12 months, fusion was achieved in 116 of 125 PEMF patients and 104 of 120 control patients (p=.1129). Patient compliance, which was automatically monitored by the device, was assessed at each visit; however, compliance data were not included in the paper. The large number of dropouts, non-significant difference in fusion rates by intent-to-treat analysis, and lack of data on functional outcomes (e.g., pain, return to usual activity) limit interpretation of these study results. Thus, this technique is considered investigational for the cervical spine.

The American Academy of Orthopedic Surgeons (AAOS) publishes information on Nonunions. “Nonunions” occur when a broken bone fails to heal and a “delayed union” is when a fracture takes longer than usual to heal. Some broken bones do not heal even when they get the best surgical or nonsurgical treatment because of inadequate stability, the blood supply is limited or lack of good nutrition to promote healing. Some bones can be expected to heal with minimal treatment due to inherent stability and excellent blood supply (toe bones). Other bones may not heal as quickly due to a limited blood supply (femoral head and neck, small wrist bone (scaphoid). Bones with moderate blood supply (tibia) may not heal quickly because the skin and muscle over the bone was damaged and the external blood supply was impaired. In addition, certain risk factors make it more likely that a bone will fail to heal. These risk factors include tobacco or nicotine use in any form, older age, severe anemia, diabetes, hypothyroidism, infection, certain medications, and low vitamin D level.

### CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

#### CPT Codes

Code	Description
20974	Electrical stimulation to aid bone healing; non-invasive (non-operative)
20975	invasive (operative)
20979	Low intensity ultrasound stimulation to aid bone healing, noninvasive (nonoperative)

Copyright © 2019 American Medical Association, Chicago, IL

#### HCPCS Codes

Code	Description
E0747	Osteogenesis stimulator, electrical, non-invasive; other than spinal applications
E0748	spinal applications
E0749	surgically implanted
E0760	Osteogenesis stimulator, low intensity ultrasound, non-invasive

#### ICD10 Codes

Code	Description
M80.00xS	Age-related osteoporosis with current pathological fracture, unspecified site, sequela
M80.021A- M80.879A	Osteoporosis with current pathological fracture, initial encounter for fracture (code range)

## Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON

Policy Number: 1.01.53

Page: 5 of 7

Code	Description
M84.30xS	Stress fracture, unspecified site, sequela
M84.38xS	Stress fracture, other site, sequela
M84.40xS	Pathological fracture, unspecified site, sequela
M84.421A- M84.48xS	Pathological fracture, initial encounter for fracture (code range)
M84.68xS	Pathological fracture in other disease, other site, sequela
Q68.8	Other specified congenital musculoskeletal deformities
Q71.61-Q71.63	Lobster-claw right hand, left hand, bilateral (code range)
Q74.0-Q74.9	Other congenital malformations of limb(s) (code range)
Q87.0	Congenital malformation syndromes predominantly affecting facial appearance
S42.201A- S42.9XS	Fracture of shoulder and upper arm, initial encounter for closed or open fracture (code range)
S49.001A- S49.199A	Other and unspecified injuries of shoulder and upper arm, initial encounter for closed fracture (code range)
S52.001A- S52.92XR	Fracture of forearm, initial encounter for closed or open fracture, initial encounter for open fracture type IIIA, IIIB, or IIIC (code range)
S59.101A- S59.199A	Other and unspecified injuries of elbow and forearm, initial encounter for closed fracture (code range)
S72.001A- S72.499C	Fracture of femur, initial encounter for closed fracture, initial encounter for open fracture type I or II, initial encounter for open fracture type IIIA, IIIB, or IIIC (code range)
S79.001A- S79.199A	Other and unspecified injuries of hip and thigh, initial encounter for closed fracture, (code range)
S82.101A- S82.866C	Fracture of lower leg, including ankle, initial encounter for closed fracture, initial encounter for open fracture type I or II, initial encounter for open fracture type IIIA, IIIB, or IIIC (code range)
S89.001A- S89.299A	Other and unspecified injuries of lower leg, initial encounter for closed fracture (code range)
S92.301A- S92.356B	Fracture of unspecified metatarsal bone(s) and great toe, initial encounter for closed or open fracture, (code range)

## REFERENCES

\*Akai M, et al. Effect of electrical stimulation on musculoskeletal systems; a meta-analysis of controlled clinical trials. Bioelectromagnetics 2002;23:132-43.

Alem IS, et al. Efficacy of electrical stimulators for bone healing: a meta-analysis of randomized sham-controlled trials. Sci Rep. Aug 19 2016;6:31724.

American Academy of Orthopedic Surgeons (AAOS). Nonunions. 2005 Mar, Updated 2007 Sep, Reviewed 2014 Mar. [<https://orthoinfo.aaos.org/en/diseases--conditions/nonunions/>] accessed 1/31/19.

BlueCross BlueShield Association. Electrical bone growth stimulation of the appendicular skeleton. Medical Policy Reference Manual Policy #7.01.07. 2018 Apr 12.

BlueCross BlueShield Association. Ultrasound accelerated fracture healing device. Medical Policy Reference Manual Policy #1.01.05. 2018 Feb 13.

\*Busse JW, et al. The effect of low-intensity pulsed ultrasound therapy on time to fracture healing: a meta-analysis. CMAJ 2002 Feb 19;166(4):437-41.

\*Busse JW, et al. Low intensity pulsed ultrasonography for fractures: systematic review of randomized controlled trials. BMJ 2009;338:b351.

## **Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON**

**Policy Number: 1.01.53**

**Page: 6 of 7**

\*Dhawan SK, et al. The effect of pulsed electromagnetic fields on hindfoot arthrodesis: a prospective study. J Foot Ankle Surg 2004 Mar-Apr;43(2):93-6.

D'Oro A, et al. Trends and costs of external electrical bone stimulators and grafting materials in anterior lumbar interbody fusion. Asian Spine J. 2018 Dec;12(6):973-980.

Ebrahim S, et al. Low-intensity pulsed ultrasonography versus electrical stimulation for fracture healing: a systematic review and network meta-analysis. Can J Surg 2014 Jun;57(3):E105-18.

\*Foley KT, et al. Randomized, prospective, and controlled clinical trial of pulsed electromagnetic field stimulation for cervical fusion. Spine J 2008;8(3):436-42.

Griffin XL, et al. Electromagnetic field stimulation for treating delayed union or non-union of long bone fractures in adults. Cochrane Database Syst Rev 2011;(4):CD008471.

Griffin XL, et al. Ultrasound and shockwave therapy for acute fractures in adults. Cochrane Database Syst Rev 2014;6:CD008579.

Hannemann PF, et al. CT scan evaluated outcome of pulsed electromagnetic fields in the treatment of acute scaphoid fractures. Bone Joint J Aug 2014;96-B(8):1070-6.

Hannemann PF, et al. The clinical and radiological outcome of pulsed electromagnetic field treatment for acute scaphoid fractures: a randomised double-blind placebo-controlled multicentre trial. J Bone Joint Surg Br 2012;94(10):1403-8.

Khalifeh JM, et al. Electrical stimulation and bone healing: a review of current technology and clinical applications. IEEE Rev Biomed Eng. 2018;11:217-232.

\*Leung KS, et al. Complex tibial fracture outcomes following treatment with low-intensity pulsed ultrasound. Ultrasound Med Biol 2004 Mar;30(3):389-95.

\*Livesley PJ, et al. Electrotherapy and the management of minimally displaced fracture of the neck of the humerus. Injury 1992;23(5):323-7.

National Institute for Health and Care Excellence. NICE medical technology guidance 12: EXOGEN ultrasound bone healing system for long bone fractures with non-union or delayed healing. 2013 Jan [http://www.nice.org.uk/guidance/mtg12] accessed 1/31/19.

Poolman RW, et al. Low intensity pulsed ultrasound (LIPUS) for bone healing: a clinical practice guideline. BMJ. Feb 21 2017;356:j576.

\*Rue JPH, et al. The effect of pulsed ultrasound in the treatment of tibial stress fractures. Orthoped 2004 Nov;27(11):1192-5.

\*Rutten S, et al. Use of low-intensity pulsed ultrasound for posttraumatic nonunions of the tibia: a review of patients treated in the Netherlands. J Trauma 2007 Apr;62:902-8.

Schandelmaier S, et al. Low intensity pulsed ultrasound for bone healing: systematic review of randomized controlled trials. BMJ. Feb 22 2017;356:j656.

Shi H, et al. Early application of pulsed electromagnetic field in the treatment of postoperative delayed union of long-bone fractures: a prospective randomized controlled study. BMC Musculoskeletal Disorders 2013;14:35.

\*Simmons JW Jr, et al. Pseudarthrosis after lumbar spine fusion: nonoperative salvage with pulsed electromagnetic fields. Am J Orthop 2004 Jan;33(1):27-30.

\*Simonis RB, et al. Electrical treatment of tibial non-union: a prospective, randomized, double-blind trial. Injury 2003 May;34(5):357-62.

\*Key Article

**Medical Policy: BONE GROWTH STIMULATORS FOR THE APPENDICULAR SKELETON**

**Policy Number: 1.01.53**

**Page: 7 of 7**

**KEY WORDS**

Bone Growth Stimulator, Osteogenic Stimulator, SAFHS, Ultrasonic Bone Growth Stimulator, US.

**CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a National Coverage Determination (NCD) for Osteogenic Stimulators. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=65&ncdver=2&bc=BAABAAAAAAAA&>