

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	ENDOMETRIAL ABLATION
Policy Number	4.01.01
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	05/18/05, 03/16/06, 03/15/07, 03/20/08, 03/19/09, 03/18/10, 04/21/11, 04/19/12, 03/21/13, 03/20/14, 03/19/15, 03/17/16, 03/16/17, 02/15/18
Archived Date	(12/20/01-05/18/05), 02/21/19
Edited Date	
Product Disclaimer	<ul style="list-style-type: none"> • <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i> • <i>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</i> • <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i>

POLICY STATEMENT

- I. Based upon our criteria and review of the peer-reviewed literature, endometrial ablation, with devices approved by the U.S. Food and Drug Administration (FDA), is **medically appropriate** and can be considered a treatment option for abnormal uterine bleed (menorrhagia) in women for whom child bearing is complete and symptoms are severe enough to warrant surgical intervention (e.g., hysterectomy).

All the following criteria must be met:

- A. Abnormal uterine bleed (menorrhagia/menometrorrhagia) for greater than 3 menstrual cycles that interferes with activities of daily living (ADLs) or results in anemia unresponsive to treatment;
- B. Pap smear in the past 12 months within normal limits;
- C. Treatment and failure to respond to hormone therapy (contraceptives, progestin) for 3 consecutive menstrual cycles or contraindication to hormone therapy;
- D. Endometrium normal within the last 6-12 months by one of the following:
 1. Hysterectomy with dilation and curettage (D & C), or
 2. Transvaginal ultrasound, or
 3. Sonohysterogram

II. Contraindications:

- A. Contraindications for endometrial ablation include:
 1. Known or suspected endometrial carcinoma or pre-malignant change of the endometrium (e.g., pre-cancerous endometrial abnormalities);
 2. Presence of enlarged uterus (e.g., greater than 10 cm in length or comparable to 12 weeks gestation or more
 3. Any anatomic or pathologic condition in which weakness of the myometrium could exist (e.g., history of previous classical cesarean section(s), transmural myomectomy);
 4. Uterine prolapse;
 5. Submucosal myomas;
 6. Active genital or urinary tract infection (e.g., cervicitis, vaginitis, endometritis, salpingitis, or cystitis);
 7. Pregnancy or desire to become pregnant in the future;
 8. Intrauterine device (IUD) in place; or
 9. Active pelvic inflammatory disease
- B. Thermal balloon endometrial ablation is contraindicated in patients who have a history of latex allergy or who have demonstrated sensitivity to latex material.
- C. Microwave ablation is contraindicated in patients who have:
 1. Essure® contraceptive micro-inserts in place,

Medical Policy: ENDOMETRIAL ABLATION

Policy Number: 4.01.01

Page: 2 of 5

2. Myometrial thickness less than 10 mm, and
 3. Uterine sounding length less than 6 cm.
- III. Based upon our criteria and review of the peer-reviewed literature, all other methods of endometrial ablation (e.g., chemoablation, photodynamic endometrial ablation) have not been medically proven to be effective and are considered **investigational**.

POLICY GUIDELINES

- I. Women with abnormal uterine bleed (menorrhagia) should be screened for possible reasons for the abnormal uterine bleed (menorrhagia) and if results appear positive further hematologic work-up should be performed. Examples of “red flags” indicating further work-up should be completed include a patient with a relative who has an inherited bleeding disorder, prolonged bleeding from small wounds or following dental procedures, heavy and prolonged bleeding following surgical procedures, easy bruising, spontaneous nosebleeds, blood in the stool or bleeding ulcer requiring urgent medical care, anemia requiring transfusion, heavy menses resulting in anemia, passing of large clots with menses or soaking more than one pad hourly, or heavy bleeding during or following childbirth.
- II. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the FDA may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Endometrial ablation is a method of treating abnormal uterine bleed (menorrhagia) through destruction of the endometrial lining. Endometrial ablation is an alternative to hysterectomy for women with abnormal uterine bleed (menorrhagia) from benign causes, who have found medical therapy ineffective or contraindicated.

In addition, in order to exclude other conditions, thyroid stimulating hormone (TSH) and human chorionic gonadotropin (HCG) testing are often performed prior to endometrial ablation to confirm these are within normal limits.

Several devices have been developed that utilize various modalities to accomplish endometrial ablation, including but not limited to: laser therapy, resecting loop rollerball using electric current, thermal ablation using a liquid-filled balloon, microwave, electrode array or a cryosurgical device.

Thermal fluid-filled balloon, cryosurgical endometrial ablation, instillation of heated saline, and radiofrequency ablation can be performed without general anesthesia in a physician’s office and do not require hysteroscopic guidance. Microwave ablation with the MEA System may also be performed in a physician’s office but does require use of the hysteroscope.

Methods that utilize direct hysteroscopic visualization include, but are not limited to:

- I. Hydrothermal (e.g. Hydro ThermAblator®, Genesys HTA™ System),
- II. Neodymium-yttrium aluminum garnet (Nd-YAG) laser,
- III. Resectoscope/resecting loop, and
- IV. Rollerball.

Methods that do not utilize direct hysteroscopic visualization include, but are not limited to:

- I. Cryoablation (e.g. Her Option™),
- II. Laser interstitial hyperthermy,
- III. Microwave (e.g. MEA System),
- IV. Radiofrequency (e.g. NovaSure™), and
- V. Thermal balloon (e.g. ThermaChoice®).

RATIONALE

Several first-generation hysteroscopically aided and second-generation non-hysteroscopically aided devices have been approved by the FDA as a safe and effective alternative to hysterectomy in select patients.

Several studies have been published addressing the various techniques of endometrial ablation as an alternative to hysterectomy for the treatment of abnormal uterine bleed (menorrhagia). In summary the studies show that endometrial

Medical Policy: ENDOMETRIAL ABLATION

Policy Number: 4.01.01

Page: 3 of 5

ablation has become the surgical treatment of choice for dysfunctional uterine bleeding when hysterectomy is not desired. In the short-term, hysteroscopic and non-hysteroscopic endometrial ablation techniques have been proven to be safe and effective in reducing excessive menstrual bleeding.

The American College of Obstetricians and Gynecologists (ACOG) practice bulletin addressing endometrial ablation states “the following recommendations and conclusions are based on good and consistent scientific evidence (Level A):

- I. For women with normal endometrial cavities, resectoscopic endometrial ablation and nonresectoscopic endometrial ablation systems appear to be equivalent with respect to successful reduction in menstrual flow and patient satisfaction at 1 year following index surgery.
- II. Resectoscopic endometrial ablation is associated with a high degree of patient satisfaction but not as high as hysterectomy.”

In the 2017 article (Klebanoff, et. al.), the authors sought to determine the incidence and predictors of failed standard of care, second-generation endometrial ablation. Failed is defined as need for surgical re-intervention. The retrospective cohort study was conducted on subjects undergoing second-generation endometrial ablation between October 2003 and March 2016. Second-generation devices utilized during the study period included the radiofrequency ablation device (RFA), hydrothermal ablation device (HTA), and the uterine balloon ablation system (UBA). Of the 5,936 women the surgical re-intervention rate was found to be 15.6%. Age, ethnicity, and radiofrequency ablation were significant risk factors for failed endometrial ablation, and menorrhagia was the leading indication for re-intervention.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
58353	Endometrial ablation, thermal, without hysteroscopic guidance
58356	Endometrial cryoablation with ultrasonic guidance, including endometrial curettage, when performed
58563	Hysteroscopy, surgical; with endometrial ablation (eg, endometrial resection, electrosurgical ablation, thermoablation)

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No specific code(s)	

ICD10 Codes

Medically Appropriate codes for when criteria are met under Policy Statement I:

Code	Description
N92.0	Excessive and frequent menstruation with regular cycle
N92.1	Excessive and frequent menstruation with irregular cycle
N92.4	Excessive bleeding in the premenopausal period
N93.8	Other specified abnormal uterine and vaginal bleeding
N93.9	Abnormal uterine and vaginal bleeding, unspecified

Medical Policy: ENDOMETRIAL ABLATION

Policy Number: 4.01.01

Page: 4 of 5

Contraindicated conditions Policy Statement II (not an all inclusive list of codes)

Code	Description
D07.0	Carcinoma in situ of endometrium
D25.0	Submucous leiomyoma of uterus
N81.2	Incomplete uterovaginal prolapse
N81.3	Complete uterovaginal prolapse
N81.4	Uterovaginal prolapse, unspecified
N85.00	Endometrial hyperplasia, unspecified
N85.02	Endometrial intraepithelial neoplasia [EIN]
N85.2	Hypertrophy of uterus

REFERENCES

American College of Obstetricians and Gynecologists. Committee Opinion 580: von Willebrand Disease in women. 2013 Dec, reaffirmed 2017.

[http://www.acog.org/Resources_And_Publications/Committee_Opinions/Committee_on_Adolescent_Health_Care/Von_Willebrand_Disease_in_Women] Accessed 1/15/19.

American College of Obstetricians and Gynecologists. Practice Bulletin #81: Endometrial ablation. *Obstet Gynecol* 2007 May;109(5):1233-48, reaffirmed 2015.

American College of Obstetricians and Gynecologists. Endometrial Ablation. FAQ 134, July 2017.

[<https://www.acog.org/Patients/FAQs/Endometrial-Ablation>] accessed 1/15/19.

*American Society for Reproductive Medicine. Indications and options for endometrial ablation. *Fertil Steril* 2008 Nov;90(5 Suppl):S236-40.

Angioni S, et al. Endometrial ablation: first- vs. second-generation techniques. *Minerva Ginecol*. Apr 2016;68(2):143-153.

Bansi-Matharu L, et al. Rates of subsequent surgery following endometrial ablation among English women with menorrhagia: population-based cohort study. *BJOG* 2013 Nov;120(12):1500-7.

Berman JM, et al. Analysis of the safety and reliability of a hydrothermal ablation system: a multicenter, prospective postmarket study. *J Reprod Med* 2014 May-Jun;59(5-6):299-305.

*Bhattacharya S, et al. Hysterectomy, endometrial ablation and Mirena® for heavy menstrual bleeding: a systematic review of clinical effectiveness and cost-effectiveness analysis. *Health Technol Assess* 2011 Apr;15(19):iii-xvi,1-252.

Blue Cross Blue Shield Association. Endometrial ablation. Medical Policy Reference Manual Policy #4.01.04. 2018 Sept 13

* Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Intrauterine ablation or resection of the endometrium for menorrhagia. TEC Evaluations 1991: Volume 6, p. 296-323.

*Daniels JP, et al; International Heavy Menstrual Bleeding IPD Meta-analysis Collaborative Group. Second generation endometrial ablation techniques for heavy menstrual bleeding: network meta-analysis. *BMJ* 2012 Apr 23;344:e2564. Daniels JP. The long-term outcomes of endometrial ablation in the treatment of heavy menstrual bleeding. *Curr Opin Obstet Gynecol* 2013 Aug;25(4):320-6.

*El-Nashar SA, et al. Global endometrial ablation for menorrhagia in women with bleeding disorders. *Obstet Gynecol* 2007 Jun;109(6):1381-7.

Fergusson RJ, et al. Endometrial resection and ablation versus hysterectomy for heavy menstrual bleeding. Cochrane Review, Cochrane Database of Systematic Reviews 2013, Issue 11, No. CD000329.

Gimpelson RJ. Ten-year literature review of global endometrial ablation with the NovaSure device *Int J Womens Health*. 2014 Mar 11;6:269-80.

Medical Policy: ENDOMETRIAL ABLATION

Policy Number: 4.01.01

Page: 5 of 5

*Hoaglin DC, et al. Use of mixed-treatment-comparison methods in estimating efficacy of treatments for heavy menstrual bleeding. Eur J Med Res 2013 Jun 21;18:17.

Klebanoff J, et al. Incidence and predictors of failed second-generation endometrial ablation. Gynecol Surg. 2017;14(1):26.

Laberge P, et al; Clinical Practice-Gynaecology Committee, Society of Obstetricians and Gynaecologists of Canada. Endometrial ablation in the management of abnormal uterine bleeding. J Obstet Gynaecol Can 2015 Apr;37(4):362-79.

Lethaby A, et al. Endometrial resection and ablation techniques for heavy menstrual bleeding. Cochrane Database of Systematic Reviews 2013, Issue 8, Art. No. CD001501.

Madsen AM, et al. Endometrial ablation for the treatment of heavy menstrual bleeding in obese women. Int J Gynaecol Obstet 2013 Apr;121(1):20-3.

Mengerink BB, et al. Effect of undiagnosed deep adenomyosis after failed NovaSure endometrial ablation. J Minim Invasive Gynecol 2015 Feb;22(2):239-44.

Miller JD, et al. Economic evaluation of global endometrial ablation versus inpatient and outpatient hysterectomy for treatment of abnormal uterine bleeding: US commercial and Medicaid payer perspectives. Popul Health Manag. 2018 Mar;21(S1):S1-S12.

Nakayama K, et al. Microwave endometrial ablation at a frequency of 2.45 GHz for menorrhagia: analysis of treatment results at a single facility. J Obstet Gynaecol Res 2014 Jan;40(1):224-9.

*NICE. Photodynamic endometrial ablation. National Institute for Clinical Excellence [Internet]. Available from <https://www.nice.org.uk/guidance/ipg47> [cited 2004, October 1]. Accessed 1/15/19.

Philip CA, et al. Evaluation of NovaSure® global endometrial ablation in symptomatic adenomyosis: a longitudinal study with a 36 month follow-up. Eur J Obstet Gynecol Reprod Biol. 2018 Aug;227:46-51.

Peeters JA, et al. Prognostic factors for the success of endometrial ablation in the treatment of menorrhagia with special reference to previous cesarean section. Eur J Obstet Gynecol Reprod Biol 2013 Mar;167(1):100-3.

Sambrook AM, et al. Microwave endometrial ablation versus thermal balloon endometrial ablation (MEATBall): 5-year follow up of a randomised controlled trial. BJOG 2014 May;121(6):747-53; discussion 754.

Thiel JA, et al. Evaluation of the NovaSure endometrial ablation procedure in women with uterine cavity length over 10 cm. J Obstet Gynaecol Can 2014 Jun;36(6):491-7.

Vilos GA, et al. Long-term clinical outcomes following resectoscopic endometrial ablation of non-atypical endometrial hyperplasia in women with abnormal uterine bleeding. J Minim Invasive Gynecol 2015 Jan;22(1):66-77.

Zupi E, et al. Hysteroscopic endometrial resection versus laparoscopic supracervical hysterectomy for abnormal uterine bleeding: long-term follow-up of a randomized trial. J Minim Invasive Gynecol 2015 Jul-Aug;22(5):841-5.

*Key Article

KEY WORDS

Endometrial ablation, Her Option™, Hydro ThermAblator®, MEA System, Novasure™, Resectoscope, Resecting loop, Rollerball, ThermaChoice®, Thermal balloon therapy.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

Based on our review, endometrial ablation is not addressed in a National or Local Medicare coverage determination or policy.