

MEDICAL POLICY



MEDICAL POLICY DETAILS	
Medical Policy Title	SURGICAL VENTRICULAR REDUCTION (e.g., PARTIAL VENTRICULECTOMY - Batista Procedure)
Policy Number	7.01.31
Category	Technology Assessment
Effective Date	10/18/01
Archived Date	10/31/01
Edited Date	09/08/05, 03/01/06, 04/19/07, 04/17/08, 04/16/09, 04/22/10, 04/21/11, 04/19/12, 04/18/13, 04/17/14, 04/16/15, 04/21/16, 04/20/17, 04/19/18, 04/18/19
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

Based upon our criteria and assessment of the peer-reviewed literature, partial ventriculectomy does not improve patient outcomes and therefore, is considered **not medically necessary** for the management of ischemic and idiopathic cardiomyopathy and as a bridge to transplant.

Refer to Corporate Medical Policy #7.01.71 regarding Surgical Ventricular Restoration.

Refer to Corporate Medical Policy # 11.01.03 Experimental and Investigational Services

POLICY GUIDELINES

The Federal Employee Health Benefit Program (FEHBP/FEP) dictates that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Patients with ischemic and idiopathic cardiomyopathy quite often have poor functioning dilated hearts. Even with maximized medical therapy and management mortality and morbidity is high. Surgical options to reduce the size of the enlarged left ventricle and improve cardiac function include partial ventriculectomy also known as the Batista procedure. Partial ventriculectomy involves removing an elliptical section of the ventricle to improve cardiac output in patients who have severe chronic heart failure. It is primarily directed at patients awaiting cardiac transplantation.

RATIONALE

There are no randomized clinical trials that support the use of surgical interventions in remodeling ventricles to treat ischemic and idiopathic cardiomyopathy. Evidence is insufficient to support surgical ventricular reduction based on small numbers of subjects and lack of availability of controlled comparisons to medical therapy, ventricular assist devices or cardiac transplantation. Additionally, the trials were unable to show that this procedure is as beneficial as other established alternatives.

In 2005, the results of the 4th International Registry Report were published, including data through 2004. This report noted that the incidence of left ventriculectomy reached a peak by 1998 and was largely abandoned by 2001, except in Asia, where experience institutions continue to perform the procedure in patients in better condition with preserved myocardial contractility.

Medical Policy: SURGICAL VENTRICULAR REDUCTION (e.g., PARTIAL VENTRICULECTOMY - Batista Procedure)

Policy Number:7.01.31

Page: 2 of 3

According to the ACC/AHA (American College of Cardiology/American Heart Association) Guideline Update for the Management of Chronic Heart Failure in the Adult (2005), although left ventriculectomy (e.g., Batista procedure) generated considerable excitement as a potential surgical approach for the treatment of refractory heart failure, the procedure failed to result in clinical improvement and was associated with a high risk of death. The ACC/AHA guideline considers ventriculectomy a Class III procedure, indicating evidence and/or general agreement that the procedure/therapy is not useful/effective and in some cases may be harmful.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
33542 (NMN)	Myocardial resection (e.g., ventricular aneurysmectomy)

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No codes	

ICD10 Codes

Code	Description
	Investigational for all diagnosis codes

REFERENCES

*Ascione R, et al. Early and late results of partial left ventriculectomy: single center experience and review of the literature. J Card Surg 2003 May-Jun;18(3):190-6.

*BlueCross BlueShield Association Technology Evaluation Center Assessment (TEC). 1998;13(4).

*BlueCross BlueShield Association. Partial left ventriculectomy. Medical Policy Reference Manual Policy #7.01.66. archived 2012 Aug 9.

Hunt SA, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to update the 2001 guidelines for the evaluation and management of heart failure). American College of Cardiology Web site [<http://content.onlinejacc.org/cgi/reprint/46/6/e1>] accessed 3/2/18.

Kawaguchi AT, et al. Left ventricular volume reduction surgery: The 4th International Registry Report 2004. J Card Surg 2005 Nov-Dec;20(6):S5-11.

*Moreira L, et al. Partial left ventriculectomy with mitral valve preservation in the treatment of patients with dilated cardiomyopathy J Thorac Cardiovasc Surg 1998;115:800-7.

National Institute for Clinical Excellence (NICE). Partial left ventriculectomy (the Batista procedure). Intervention procedure guidance 41. NICE 2004 Feb [<https://www.nice.org.uk/guidance/ipg41>] accessed 3/2/18.

*Popovic Z, et al. Effects of partial left ventriculectomy on left ventricular performance in patients with nonischemic dilated cardiomyopathy J Am Coll Cardiol 1998;32:1801-8.

Medical Policy: SURGICAL VENTRICULAR REDUCTION (e.g., PARTIAL VENTRICULECTOMY - Batista Procedure)

Policy Number: 7.01.31

Page: 3 of 3

*Stolf N, et al. Determinants of midterm outcome of partial left ventriculectomy in dilated cardiomyopathy Ann Thorac Surg 1998;66:1585-91.

Tonnessen T, et al. Surgical left ventricular remodeling in heart failure. Eur J Heart Fail 2005 Aug;7(5):704-9.

*key articles

KEY WORDS

Cardio-reduction, ventriculectomy, ventricular remodeling, Batista procedure

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for partial ventriculectomy. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=122&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&CptHcpcsCode=36514&bc=gAAAABAAAA&>.